



planetmath.org

Math for the people, by the people.

anti-isomorphism

Canonical name	Antiisomorphism
Date of creation	2013-03-22 16:01:08
Last modified on	2013-03-22 16:01:08
Owner	Mathprof (13753)
Last modified by	Mathprof (13753)
Numerical id	15
Author	Mathprof (13753)
Entry type	Definition
Classification	msc 13B10
Classification	msc 16B99
Defines	anti-endomorphism
Defines	anti-homomorphism
Defines	anti-isomorphic
Defines	anti-automorphism

Let R and S be rings and $f : R \longrightarrow S$ be a function such that $f(r_1r_2) = f(r_2)f(r_1)$ for all $r_1, r_2 \in R$.

If f is a homomorphism of the additive groups of R and S , then f is called an *anti-homomorphism*.

If f is a bijection and anti-homomorphism, then f is called an *anti-isomorphism*.

If f is an anti-homomorphism and $R = S$ then f is called an *anti-automorphism*.

If f is an anti-isomorphism and $R = S$ then f is called an *anti-automorphism*.

As an example, when $m \neq n$, the mapping that sends a matrix to its transpose (or to its conjugate transpose if the matrix is complex) is an anti-isomorphism of $M_{m,n} \rightarrow M_{n,m}$.

R and S are *anti-isomorphic* if there is an anti-isomorphism $R \rightarrow S$.

All of the things defined in this entry are also defined for groups.