



Math for the people, by the people.

### applying elementary symmetric polynomials

Canonical name	ApplyingElementarySymmetricPolynomials
Date of creation	2013-03-22 19:10:07
Last modified on	2013-03-22 19:10:07
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	16
Author	pahio (2872)
Entry type	Application
Classification	msc 13B25
Classification	msc 12E10

The method used in the proof of fundamental theorem of symmetric polynomials may be applied to concrete instances as follows.

We assume the given a symmetric polynomial  $P(x_1, x_2, \dots, x_n) = P$  of degree  $d$  be <http://planetmath.org/HomogeneousPolynomialhomogeneous>. Starting from the highest term of  $P$  we form all products

$$x_1^{\lambda_1} x_2^{\lambda_2} \dots x_n^{\lambda_n}$$

where

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0 \quad \text{and} \quad \lambda_1 + \lambda_2 + \dots + \lambda_n = d.$$

Then

$$P = Q(p_1, p_2, \dots, p_n) = \sum_i m_i p_1^{\lambda_1 - \lambda_2} p_2^{\lambda_2 - \lambda_3} \dots p_{n-1}^{\lambda_{n-1} - \lambda_n} p_n^{\lambda_n}, \quad (1)$$

in which the coefficients  $m_i$  are determined by giving some suitable values to the indeterminates  $x_j$ .

**Example 1.** Express the polynomial  $P = x_1^3 x_2 + x_1^3 x_3 + x_2^3 x_1 + x_2^3 x_3 + x_3^3 x_1 + x_3^3 x_2$  in the elementary symmetric polynomials

$$p_1 = x_1 + x_2 + x_3, \quad p_2 = x_2 x_3 + x_3 x_1 + x_1 x_2, \quad p_3 = x_1 x_2 x_3. \quad (2)$$

We have four

$$4, 0, 0; \quad 3, 1, 0; \quad 2, 2, 0; \quad 2, 1, 1,$$

for which the corresponding  $p$ -products of the sum (1) are

$$p_1^4, \quad p_1^2 p_2, \quad p_2^2, \quad p_1 p_3,$$

respectively. Apparently, the first one is out of the question. Therefore, clearly

$$P = p_1^2 p_2 + a p_2^2 + b p_1 p_3.$$

Using  $x_1 = x_2 = 1$  and  $x_3 = 0$  makes  $p_1 = 2$ ,  $p_2 = 1$  and  $p_3 = 0$ , when

$$P = 2 = 4 + a + 0,$$

implying  $a = -2$ . Using similarly  $x_1 = x_2 = x_3 = 1$  we get  $p_1 = p_2 = 3$ ,  $p_3 = 1$ , which give

$$P = 6 = 27 + 9a + 3b = 9 + 3b,$$

yielding  $b = -1$ . Hence we have the result

$$P = p_1^2 p_2 - 2p_2^2 - p_1 p_3,$$

i.e.

$$x_1^3 x_2 + x_1^3 x_3 + x_2^3 x_1 + x_2^3 x_3 + x_3^3 x_1 + x_3^3 x_2 = (x_1 + x_2 + x_3)^2 (x_2 x_3 + x_3 x_1 + x_1 x_2) - 2(x_2 x_3 + x_3 x_1 + x_1 x_2)^2 - (x_1 + x_2 + x_3)$$

**Example 2.** Let  $P = x_1^4 + x_2^4 + \dots + x_n^4$ . If we suppose that  $n \geq 4$ , the possible highest terms are

$$x_1^4, \quad x_1^3 x_2, \quad x_1^2 x_2^2, \quad x_1^2 x_2 x_3, \quad x_1 x_2 x_3 x_4$$

whence we may write

$$P = p_1^4 + a p_1^2 p_2 + b p_2^2 + c p_1 p_3 + d p_4. \quad (3)$$

For determining the coefficients, evidently we can put  $x_5 = x_6 = \dots = x_n = 0$  and in as follows.

1°.  $x_1 = 1, x_2 = -1, x_3 = x_4 = 0$ . Then we have  $P = 2, p_1 = 0, p_2 = -1, p_3 = p_4 = 0$ . Thus (3) gives  $b = 2$ .

2°.  $x_1 = x_2 = 1, x_3 = x_4 = -1$ . Now  $P = 4, p_1 = 0, p_2 = -2, p_3 = 0, p_4 = 1$ , whence (3) reads  $4 = 4b + d = 8 + d$ , giving  $d = -4$ .

3°.  $x_1 = x_2 = 1, x_3 = x_4 = 0$ . We get  $P = 2, p_1 = 2, p_2 = 1, p_3 = p_4 = 0$ . These yield  $2 = 16 + 4a + b = 18 + 4a$ , i.e.  $a = -4$ .

4°.  $x_1 = x_2 = 2, x_3 = -1, x_4 = 0$ . In this case,  $P = 33, p_1 = 3, p_2 = 0, p_3 = -4, p_4 = 0$ , whence  $33 = 81 - 12c$ , or  $c = 4$ . Consequently, we obtain from (3) the result

$$P = p_1^4 - 4p_1^2 p_2 + 2p_2^2 + 4p_1 p_3 - 4p_4. \quad (4)$$

Although it has been derived by supposing  $n \geq 4$  (= the degree of  $P$ ), it holds without this supposition. One has only to see that e.g. in the case  $n = 2$ , one must substitute to (4) the values  $p_3 = p_4 = 0$ , which changes the to the form  $P = p_1^4 - 4p_1^2 p_2 + 2p_2^2$ .