



planetmath.org

Math for the people, by the people.

continuation of exponent

Canonical name	ContinuationOfExponent
Date of creation	2013-03-22 17:59:49
Last modified on	2013-03-22 17:59:49
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	6
Author	pahio (2872)
Entry type	Definition
Classification	msc 13A18
Classification	msc 12J20
Classification	msc 11R99
Classification	msc 13F30
Synonym	prolongation of exponent
Defines	induce
Defines	continuation
Defines	continuation of the exponent
Defines	ramification index of the exponent

Theorem. Let K/k be a finite field extension and ν an exponent valuation of the extension field K . Then there exists one and only one positive integer e such that the function

$$(1) \quad \nu_0(x) := \begin{cases} \infty & \text{when } x = 0, \\ \frac{\nu(x)}{e} & \text{when } x \neq 0, \end{cases}$$

defined in the base field k , is an <http://planetmath.org/ExponentValuationexponent> of k .

Proof. The exponent ν of K attains in the set $k \setminus \{0\}$ also non-zero values; otherwise k would be included in \mathcal{O}_ν , the ring of the exponent ν . Since any element ξ of K are integral over k , it would then be also integral over \mathcal{O}_ν , which is integrally closed in its quotient field K (see theorem 1 in ring of exponent); the situation would mean that $\xi \in \mathcal{O}_\nu$ and thus the whole K would be contained in \mathcal{O}_ν . This is impossible, because an exponent of K attains also negative values. So we infer that ν does not vanish in the whole $k \setminus \{0\}$. Furthermore, ν attains in $k \setminus \{0\}$ both negative and positive values, since $\nu(a) + \nu(a^{-1}) = \nu(aa^{-1}) = \nu(1) = 0$.

Let p be such an element of k on which ν attains as its value the least possible positive integer e in the field k and let a be an arbitrary non-zero element of k . If

$$\nu(a) = m = qe + r \quad (q, r \in \mathbb{Z}, \quad 0 \leq r < e),$$

then $\nu(ap^{-q}) = m - qe = r$, and thus $r = 0$ on grounds of the choice of p . This means that $\nu(a)$ is always divisible by e , i.e. that the values of the function ν_0 in $k \setminus \{0\}$ are integers. Because $\nu_0(p) = 1$ and $\nu_0(p^l) = l$, the function attains in k every integer value. Also the conditions

$$\nu_0(ab) = \nu_0(a) + \nu_0(b), \quad \nu_0(a + b) \geq \min\{\nu_0(a), \nu_0(b)\}$$

are in , whence ν_0 is an exponent of the field k .

Definition. Let K/k be a finite field extension. If the exponent ν_0 of k is tied with the exponent ν of K via the condition (1), one says that ν induces ν_0 to k and that ν is the *continuation* of ν_0 to K . The positive integer e , uniquely determined by (1), is the *ramification index* of ν with respect to ν_0 (or with respect to the subfield k).

References

- [1] S. BOREWICZ & I. SAFAREVIC: *Zahlentheorie*. Birkhäuser Verlag. Basel und Stuttgart (1966).