



planetmath.org

Math for the people, by the people.

Krull valuation domain

Canonical name	KrullValuationDomain
Date of creation	2013-03-22 14:55:01
Last modified on	2013-03-22 14:55:01
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	8
Author	pahio (2872)
Entry type	Theorem
Classification	msc 13F30
Classification	msc 13A18
Classification	msc 12J20
Classification	msc 11R99
Related topic	ValuationDeterminedByValuationDomain

Theorem. Any Krull valuation $|\cdot|$ of a field K determines a unique valuation domain $R = \{a \in K : |a| \leq 1\}$, whose field of fraction is K .

Proof. We first see that $1 \in R$ since $|1| = 1$. Let then a, b be any two elements of R . The non-archimedean triangle inequality shows that $|a - b| \leq \max\{|a|, |b|\} \leq 1$, i.e. that the difference $a - b$ belongs to R . Using the <http://planetmath.org/OrderedGroupmultiplication> rule 4 of inequalities we obtain

$$|ab| = |a| \cdot |b| \leq 1 \cdot 1 = 1$$

which shows that also the product ab is element of R . Thus, R is a subring of the field K , and so an integral domain. Let now c be an arbitrary element of K not belonging to R . This implies that $1 < |c|$, whence $|c^{-1}| = |c|^{-1} < 1$ (see the <http://planetmath.org/OrderedGroupinverse> rule 5). Consequently, the inverse c^{-1} belongs to R , and we conclude that R is a valuation domain. The $a = \frac{a}{1}$ and $c = \frac{1}{c^{-1}}$ make evident that K is the field of fractions of R .