



planetmath.org

Math for the people, by the people.

## product of finitely generated ideals

Canonical name	ProductOffinitelyGeneratedIdeals
Date of creation	2015-05-05 19:19:39
Last modified on	2015-05-05 19:19:39
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	29
Author	pahio (2872)
Entry type	Definition
Classification	msc 13A15
Classification	msc 16D25
Classification	msc 16D10
Synonym	special cases of ideal product
Related topic	PruferRing
Related topic	IdealGeneratorsInPruferRing
Related topic	IdealDecompositionInDedekindDomain
Related topic	EntriesOnFinitelyGeneratedIdeals
Related topic	UniqueFactorizationAndIdealsInRingOfIntegers
Related topic	ContentOfAPolynomial
Related topic	WellDefinednessOfProductOffinitelyGeneratedIdeals
Defines	Dedekind–Mertens lemma

Let  $R$  be a commutative ring having at least one regular element and  $T$  its total ring of fractions. Let  $\mathfrak{a} := (a_0, a_1, \dots, a_{m-1})$  and  $\mathfrak{b} := (b_0, b_1, \dots, b_{n-1})$  be two fractional ideals of  $R$  (see the entry “fractional ideal of commutative ring”). Then the product submodule  $\mathfrak{a}\mathfrak{b}$  of  $T$  is also a of  $R$  and is generated by all the elements  $a_i b_j$ , thus having a generating set of  $mn$  elements.

Such a generating set may be condensed in the case of any Dedekind domain, especially for the of any algebraic number field one has the multiplication formula

$$\mathfrak{a}\mathfrak{b} = (a_0 b_0, a_0 b_1 + a_1 b_0, a_0 b_2 + a_1 b_1 + a_2 b_0, \dots, a_{m-1} b_{n-1}). \quad (1)$$

Here, the number of generators is only  $m+n-1$  (in principle, every ideal of a Dedekind domain has a <http://planetmath.org/TwoGeneratorProperty> generating system of two elements). The formula is <http://planetmath.org/Characterizationcharacteristi> still for a wider class of rings  $R$  which may contain zero divisors, viz. for the Prüfer rings (see [1]), but then at least one of  $\mathfrak{a}$  and  $\mathfrak{b}$  must be a regular ideal.

Note that the generators in (1) are formed similarly as the coefficients in the product of the polynomials  $f(X) := f_0 + f_1 X + \dots + f_{m-1} X^{m-1}$  and  $g(X) := g_0 + g_1 X + \dots + g_{n-1} X^{n-1}$ . Thus we may call the fractional ideals  $\mathfrak{a}$  and  $\mathfrak{b}$  of  $R$  the *coefficient modules*  $\mathfrak{m}_f$  and  $\mathfrak{m}_g$  of the polynomials  $f$  and  $g$  (they are  $R$ -modules). Hence the formula (1) may be rewritten as

$$\mathfrak{m}_f \mathfrak{m}_g = \mathfrak{m}_{fg}. \quad (2)$$

This formula says the same as Gauss’s lemma I for a unique factorization domain  $R$ .

Arnold and Gilmer [2] have presented and proved the following generalisation of (2) which is valid under much less stringent assumptions than the ones requiring  $R$  to be a Prüfer ring (initially: a Prüfer domain); the proof is somewhat simplified in [1].

**Theorem (Dedekind–Mertens lemma).** Let  $R$  be a subring of a commutative ring  $T$ . If  $f$  and  $g$  are two arbitrary polynomials in the polynomial ring  $T[X]$ , then there exists a non-negative integer  $n$  such that the  $R$ -submodules of  $T$  generated by the coefficients of the polynomials  $f$ ,  $g$  and  $fg$  satisfy the equality

$$\mathfrak{m}_f^{n+1} \mathfrak{m}_g = \mathfrak{m}_f^n \mathfrak{m}_{fg}. \quad (3)$$

## References

- [1] J. PAHIKKALA: “Some formulae for multiplying and inverting ideals”. – *Ann. Univ. Turkuensis* **183** (A) (1982).
- [2] J. ARNOLD & R. GILMER: “On the contents of polynomials”. – *Proc. Amer. Math. Soc.* **24** (1970).