

## planetmath.org

Math for the people, by the people.

## henselian field

Canonical name HenselianField

Date of creation 2013-03-22 13:28:37 Last modified on 2013-03-22 13:28:37

Owner mps (409) Last modified by mps (409)

Numerical id 9

Author mps (409)
Entry type Definition
Classification msc 13F30
Classification msc 13A18
Classification msc 11R99
Classification msc 12J20
Related topic Valuation

Related topic ValuationDomainIsLocal Related topic ValuationRingOfAField

Defines valuation ring
Defines residue field
Defines residue class field
Defines Hensel property

Defines henselian
Defines henselisation

Let  $|\cdot|$  be a non-archimedean valuation on a field K. Let  $V = \{x : |x| \le 1\}$ . Since  $|\cdot|$  is ultrametric, V is closed under addition and in fact an additive group. The other valuation axioms ensure that V is a ring. We call V the valuation ring of K with respect to the valuation  $|\cdot|$ . Note that the field of fractions of V is K.

The set  $\mu = \{x : |x| < 1\}$  is a maximal ideal of V. The factor  $R := V/\mu$  is called the *residue field* or the *residue class field*.

The map res :  $V \to V/\mu$  given by  $x \mapsto x + \mu$  is called the *residue map*. We extend the definition of the residue map to sequences of elements from V, and hence to V[X] so that if  $f(X) \in V[X]$  is given by  $\sum_{i \le n} a_i X^i$  then  $\operatorname{res}(f) \in R[X]$  is given by  $\sum_{i \le n} \operatorname{res}(ai) X^i$ .

**Hensel property:** Let  $f(x) \in V[x]$ . Suppose  $\operatorname{res}(f)(x)$  has a simple root  $e \in k$ . Then f(x) has a root  $e' \in V$  and  $\operatorname{res}(e') = e$ .

Any valued field satisfying the Hensel property is called *henselian*. The completion of a non-archimedean valued field K with respect to the valuation (cf. constructing the reals from the rationals as the completion with respect to the standard metric) is a henselian field.

Every non-archimedean valued field K has a unique (up to isomorphism) smallest henselian field  $K^h$  containing it. We call  $K^h$  the henselisation of K.