



planetmath.org

Math for the people, by the people.

equivalent valuations

Canonical name	EquivalentValuations
Date of creation	2013-03-22 14:25:27
Last modified on	2013-03-22 14:25:27
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	18
Author	pahio (2872)
Entry type	Definition
Classification	msc 13A18
Related topic	DiscreteValuation
Related topic	IndependenceOfTheValuations
Defines	equivalence of valuations

Let  $K$  be a field. The *equivalence of valuations*  $|\cdot|_1$  and  $|\cdot|_2$  of  $K$  may be defined so that

1.  $|\cdot|_1$  is not the trivial valuation;
2. if  $|a|_1 < 1$  then  $|a|_2 < 1 \quad \forall a \in K$ .

It is easy to see that these conditions imply for both valuations (use  $\frac{1}{a}$ ). Also, we have always

$$|a|_1 \leq 1 \Leftrightarrow |a|_2 \leq 1;$$

so both valuations have a common valuation ring in the case they are non-archimedean. (The more general Krull valuation is defined so that they have common valuation rings.) Further, both valuations determine a common metric on  $K$ .

**Theorem.** Two valuations (of <http://planetmath.org/KrullValuationrank> one)  $|\cdot|_1$  and  $|\cdot|_2$  of  $K$  are iff one of them is a positive power of the other,

$$|a|_1 = |a|_2^c \quad \forall a \in K,$$

where  $c$  is a positive .