



planetmath.org

Math for the people, by the people.

**proof that a domain is Dedekind if its ideals
are products of primes**

Canonical name	ProofThatADomainIsDedekindIfItsIdealsAreProductsOfPrimes
Date of creation	2013-03-22 18:35:07
Last modified on	2013-03-22 18:35:07
Owner	gel (22282)
Last modified by	gel (22282)
Numerical id	5
Author	gel (22282)
Entry type	Proof
Classification	msc 13A15
Classification	msc 13F05
Related topic	DedekindDomain
Related topic	PrimeIdeal

We show that for an integral domain R , the following are equivalent.

1. R is a Dedekind domain.
2. every nonzero proper ideal is a product of maximal ideals.
3. every nonzero proper ideal is a product of prime ideals.

For the equivalence of ?? and ?? see proof that a domain is Dedekind if its ideals are products of maximals. Also, as every maximal ideal is prime, it is immediate that ?? implies ?. So, we just need to consider the case where ?? is satisfied and show that ?? follows, for which it enough to show that every nonzero prime ideal is maximal.

We first suppose that \mathfrak{p} is an <http://planetmath.org/FractionalIdealinvertible> prime ideal and show that it is maximal. To do this it is enough to show that any $a \in R \setminus \mathfrak{p}$ gives $\mathfrak{p} + (a) = R$. First, we have the following inclusions,

$$\mathfrak{p} \subsetneq \mathfrak{p} + (a^2) \subseteq \mathfrak{p} + (a).$$

Then, consider the prime factorizations

$$\mathfrak{p} + (a) = \mathfrak{p}_1 \cdots \mathfrak{p}_m, \quad (1)$$

$$\mathfrak{p} + (a^2) = \mathfrak{q}_1 \cdots \mathfrak{q}_n. \quad (2)$$

We write \bar{a} for the image of a under the <http://planetmath.org/NaturalHomomorphismnatural> homomorphism $R \rightarrow R/\mathfrak{p}$ and $\bar{\mathfrak{a}}$ for the image of any ideal \mathfrak{a} . Equations (??) and (??) give

$$\bar{\mathfrak{q}}_1 \cdots \bar{\mathfrak{q}}_n = (\bar{a})^2 = (\bar{\mathfrak{p}}_1 \cdots \bar{\mathfrak{p}}_m)^2. \quad (3)$$

As \mathfrak{p} is strictly contained in $\mathfrak{p} + (a)$ and $\mathfrak{p} + (a^2)$, it must also be strictly contained in \mathfrak{p}_k and \mathfrak{q}_k . So, $\bar{\mathfrak{p}}_k, \bar{\mathfrak{q}}_k$ are nonzero prime ideals, and by uniqueness of prime factorization (see, prime ideal factorization is unique) Equation (??) gives $n = 2m$ and $\bar{\mathfrak{p}}_k = \bar{\mathfrak{q}}_k = \bar{\mathfrak{q}}_{k+m}$, after reordering of the factors. So $\mathfrak{p}_k = \mathfrak{q}_k = \mathfrak{q}_{k+m}$ and,

$$\mathfrak{p} + (a^2) = \mathfrak{q}_1 \cdots \mathfrak{q}_n = (\mathfrak{p}_1 \cdots \mathfrak{p}_m)^2 = (\mathfrak{p} + (a))^2 = \mathfrak{p}^2 + a\mathfrak{p} + (a^2). \quad (4)$$

Then, $a \notin \mathfrak{p}$ gives $\mathfrak{p} \cap (a^2) \subseteq a\mathfrak{p}$ and taking the intersection of both sides of (??) with \mathfrak{p} ,

$$\mathfrak{p} = \mathfrak{p}^2 + a\mathfrak{p} + \mathfrak{p} \cap (a^2) \subseteq \mathfrak{p}^2 + a\mathfrak{p} = (\mathfrak{p} + (a))\mathfrak{p}.$$

But \mathfrak{p} was assumed to be invertible, and can be cancelled giving $R \subseteq \mathfrak{p} + (a)$, showing that \mathfrak{p} is maximal.

Now let \mathfrak{p} be any prime ideal and $a \in \mathfrak{p} \setminus \{0\}$. Factoring into a product of primes

$$\mathfrak{p}_1 \cdots \mathfrak{p}_n = (a) \subseteq \mathfrak{p}, \tag{5}$$

each of the \mathfrak{p}_k is invertible and, by the above argument, must be maximal. Finally, as \mathfrak{p} is prime, (??) gives $\mathfrak{p}_k \subseteq \mathfrak{p}$ for some k , so $\mathfrak{p} = \mathfrak{p}_k$ is maximal.