



integral closures in separable extensions are
finitely generated

Canonical name	IntegralClosuresInSeparableExtensionsAreFinitelyGenerated
Date of creation	2013-03-22 17:02:12
Last modified on	2013-03-22 17:02:12
Owner	rm50 (10146)
Last modified by	rm50 (10146)
Numerical id	5
Author	rm50 (10146)
Entry type	Theorem
Classification	msc 13B21
Classification	msc 12F05
Related topic	IntegralClosureIsRing

The theorem below generalizes to arbitrary integral ring extensions (under certain conditions) the fact that the ring of integers of a number field is finitely generated over \mathbb{Z} . The proof parallels the proof of the number field result.

Theorem 1. *Let B be an integrally closed Noetherian domain with field of fractions K . Let L be a finite separable extension of K , and let A be the integral closure of B in L . Then A is a finitely generated B -module.*

Proof. We first show that the <http://planetmath.org/Trace2trace> Tr_K^L maps A to B . Choose $u \in A$ and let $f = Irr(u, K) \in K[x]$ be the minimal polynomial for u over K ; assume f is of degree d . Let the conjugates of u in some splitting field be $u = a_1, \dots, a_d$. Then the a_i are all integral over B since they satisfy u 's monic polynomial in $B[x]$. Since the coefficients of F are polynomials in the a_i , they too are integral over B . But the coefficients are in K , and B is integrally closed (in K), so the coefficients are in B . But $Tr_K^L(u)$ is just the coefficient of x^{d-1} in f , and thus $Tr_K^L(u) \in B$. This proves the claim.

Now, choose a basis $\omega_1, \dots, \omega_d$ of L/K . We may assume $\omega_i \in A$ by multiplying each by an appropriate element of B . (To see this, let $Irr(\omega_i, K) \in K[x] = x^d + k_1x^{d-1} + \dots + k_d$. Choose $b \in B$ such that $bk_i \in B \forall i$. Then $(b\omega)^d + bk_1(b\omega)^{d-1} + \dots + b^dk_d = 0$ and thus $b\omega \in A$). Define a linear map $\varphi : L \rightarrow K^d : a \mapsto (Tr_K^L(a\omega_1), \dots, Tr_K^L(a\omega_d))$.

φ is 1-1, since if $u \in \ker \varphi, u \neq 0$, then $Tr(uL) = 0$. But $uL = L$, so Tr_K^L is identically zero, which cannot be since L is separable over K (it is a standard result that separability is equivalent to nonvanishing of the trace map; see for example [?], Chapter 8).

But $Tr_K^L : A \rightarrow B$ by the above, so $\varphi : A \hookrightarrow B^d$. Since B is Noetherian, any submodule of a finitely generated module is also finitely generated, so A is finitely generated as a B -module. \square

References

- [1] P. Morandi, *Field and Galois Theory*, Springer, 2006.