

planetmath.org

Math for the people, by the people.

$\begin{array}{c} \textbf{Dedekind domains with finitely many primes} \\ \textbf{are PIDs} \end{array}$

 ${\bf Canonical\ name} \quad {\bf Dedekind Domains With Finitely Many Primes Are PIDs}$

Date of creation 2013-03-22 18:35:18 Last modified on 2013-03-22 18:35:18

Owner gel (22282) Last modified by gel (22282)

Numerical id 6

Author gel (22282)
Entry type Theorem
Classification msc 13F05
Classification msc 11R04
Related topic DivisorTheory

A commutative ring in which there are only finitely many maximal ideals is known as a semi-local ring. For such rings, the property of being a Dedekind domain and of being a principal ideal domain coincide.

Theorem. A Dedekind domain in which there are only finitely many prime ideals is a principal ideal domain.

This result is sometimes proven using the chinese remainder theorem or, alternatively, it follows directly from the fact that invertible ideals in semi-local rings are principal.

Suppose that R is a Dedekind domain such as the ring of algebraic integers in a number field. Although there are infinitely many prime ideals in such a ring, we can use the result that localizations of Dedekind domains are Dedekind and apply the above theorem to localizations of R.

In particular, if \mathfrak{p} is a nonzero prime ideal, then $R_{\mathfrak{p}} \equiv (R \setminus \mathfrak{p})^{-1}R$ is a Dedekind domain with a unique nonzero prime ideal, so the theorem shows that it is a principal ideal domain.