



planetmath.org

Math for the people, by the people.

height function

Canonical name	HeightFunction
Date of creation	2013-03-22 13:49:09
Last modified on	2013-03-22 13:49:09
Owner	alozano (2414)
Last modified by	alozano (2414)
Numerical id	5
Author	alozano (2414)
Entry type	Definition
Classification	msc 14H52
Related topic	EllipticCurve
Related topic	RankOfAnEllipticCurve
Related topic	ArithmeticOfEllipticCurves
Related topic	CanonicalHeightOnAnEllipticCurve
Defines	height function
Defines	canonical height
Defines	descent theorem

Definition 1 Let A be an abelian group. A height function on A is a function $h: A \rightarrow \mathbb{R}$ with the properties:

1. For all $Q \in A$ there exists a constant C_1 , depending on A and Q , such that for all $P \in A$:

$$h(P + Q) \leq 2h(P) + C_1$$

2. There exists an integer $m \geq 2$ and a constant C_2 , depending on A , such that for all $P \in A$:

$$h(mP) \geq m^2h(P) - C_2$$

3. For all $C_3 \in \mathbb{R}$, the following set is finite:

$$\{P \in A : h(P) \leq C_3\}$$

Examples:

1. For $t = p/q \in \mathbb{Q}$, a fraction in lower terms, define $H(t) = \max\{|p|, |q|\}$. Even though this is not a height function as defined above, this is the prototype of what a height function should look like.
2. Let E be an elliptic curve over \mathbb{Q} . The function on $E(\mathbb{Q})$, the points in E with coordinates in \mathbb{Q} , $h_x: E(\mathbb{Q}) \rightarrow \mathbb{R}$:

$$h_x(P) = \begin{cases} \log H(x(P)), & \text{if } P \neq 0 \\ 0, & \text{if } P = 0 \end{cases}$$

is a height function (H is defined as above). Notice that this depends on the chosen Weierstrass model of the curve.

3. The *canonical height* of E/\mathbb{Q} (due to Neron and Tate) is defined by:

$$h_C(P) = 1/2 \lim_{N \rightarrow \infty} 4^{(-N)} h_x([2^N]P)$$

where h_x is defined as in (2).

Finally we mention the fundamental theorem of “descent”, which highlights the importance of the height functions:

Theorem 1 (Descent) Let A be an abelian group and let $h: A \rightarrow \mathbb{R}$ be a height function. Suppose that for the integer m , as in property (2) of height, the quotient group A/mA is finite. Then A is finitely generated.

References

- [1] Joseph H. Silverman, *The Arithmetic of Elliptic Curves*. Springer-Verlag, New York, 1986.