



planetmath.org

Math for the people, by the people.

Shioda-Tate formula

Canonical name	ShiodaTateFormula
Date of creation	2013-03-22 15:34:22
Last modified on	2013-03-22 15:34:22
Owner	alozano (2414)
Last modified by	alozano (2414)
Numerical id	4
Author	alozano (2414)
Entry type	Theorem
Classification	msc 14J27

The main references for this part are the works of Shioda and Tate [?], [?], [?].

Let k be a field and let \bar{k} be a fixed algebraic closure of k . Let \mathcal{E} be an elliptic surface over a curve C/k and let $K = k(C)$ be the function field of C . Let $\bar{\mathcal{E}} = \mathcal{E}(\bar{k})$ (or more precisely $\bar{\mathcal{E}} = \mathcal{E} \times_{\text{Spec } k} \text{Spec } \bar{k}$). The Néron-Severi group of $\bar{\mathcal{E}}$, denoted by $\text{NS}(\bar{\mathcal{E}})$, is by definition the group of divisors on $\bar{\mathcal{E}}$ modulo algebraic equivalence. Under the previous assumptions, $\text{NS}(\bar{\mathcal{E}})$ is a finitely generated abelian group (this is a consequence of the so-called ‘theorem of the base’ which can be found in [?]). The Néron-Severi group of \mathcal{E} , denoted by $\text{NS}(\mathcal{E})$, is simply the image of the group of divisors on \mathcal{E} in $\text{NS}(\bar{\mathcal{E}})$. Let $T \subset \text{NS}(\mathcal{E})$ be the subgroup generated by the image of the zero-section σ_0 and all the irreducible components of the fibers of π . T is sometimes called the “trivial part” of $\text{NS}(\mathcal{E})$.

Theorem (Shioda-Tate formula). *For each $t \in C$ let n_t be the number of irreducible components on the fiber at t , i.e. $\pi^{-1}(t)$. Then:*

$$\begin{aligned} \text{rank}_{\mathbb{Z}}(\mathcal{E}/K) &= \text{rank}_{\mathbb{Z}}(\text{NS}(\mathcal{E})) - \text{rank}_{\mathbb{Z}}(T) \\ &= \text{rank}_{\mathbb{Z}}(\text{NS}(\mathcal{E})) - 2 - \sum_{t \in C} (n_t - 1). \end{aligned}$$

References

- [1] S. Lang, *Fundamentals of Diophantine Geometry*, Springer-Verlag (1983).
- [2] T. Shioda, *On elliptic modular surfaces*, J. Math. Soc. Japan 24 (1972), 20-59.
- [3] T. Shioda, *An Explicit Algorithm for Computing the Picard Number of Certain Algebraic Surfaces*, Amer. J. Math. 108 (1986), 415-432.
- [4] T. Shioda, *On the Mordell-Weil Lattices*, Commentarii Mathematici Universitatis Sancti Pauli, Vol 39, No. 2, 1990, pp. 211-239.
- [5] J. Tate, *On the conjectures of Birch and Swinnerton-Dyer and a geometric analog*, Séminaire Bourbaki, 9, Soc. Math. France, Paris, 1966, Exp. No. 306, 415-440, 1995.