



sheaf cohomology

Canonical name	SheafCohomology
Date of creation	2013-03-22 13:50:59
Last modified on	2013-03-22 13:50:59
Owner	mathcam (2727)
Last modified by	mathcam (2727)
Numerical id	14
Author	mathcam (2727)
Entry type	Definition
Classification	msc 14F25
Related topic	EtaleCohomology
Related topic	LeraysTheorem
Related topic	AcyclicSheaf
Related topic	DeRhamWeilTheorem
Defines	sufficiently fine

Let X be a topological space. The category of sheaves of abelian groups on X has enough injectives. So we can define the sheaf cohomology $H^i(X, \mathcal{F})$ of a sheaf \mathcal{F} to be the right derived functors of the global sections functor $\mathcal{F} \rightarrow \Gamma(X, \mathcal{F})$.

Usually we are interested in the case where X is a scheme, and \mathcal{F} is a coherent sheaf. In this case, it does not matter if we take the derived functors in the category of sheaves of abelian groups or coherent sheaves.

Sheaf cohomology can be explicitly calculated using <http://planetmath.org/CechCohomology> cohomology. Choose an open cover $\{U_i\}$ of X . We define

$$C^i(\mathcal{F}) = \prod \mathcal{F}(U_{j_0 \dots j_i})$$

where the product is over $i + 1$ element subsets of $\{1, \dots, n\}$ and $U_{j_0 \dots j_i} = U_{j_0} \cap \dots \cap U_{j_i}$. If $s \in \mathcal{F}(U_{j_0 \dots j_i})$ is thought of as an element of $C^i(\mathcal{F})$, then the differential

$$\partial(s) = \prod_{\ell} \left(\prod_{k=j_{\ell}+1}^{j_{\ell+1}-1} (-1)^{\ell} s|_{U_{j_0 \dots j_{\ell} k j_{\ell+1} \dots j_i}} \right)$$

makes $C^*(\mathcal{F})$ into a chain complex. The cohomology of this complex is denoted $\check{H}^i(X, \mathcal{F})$ and called the *Čech cohomology* of \mathcal{F} with respect to the cover $\{U_i\}$. There is a natural map $H^i(X, \mathcal{F}) \rightarrow \check{H}^i(X, \mathcal{F})$ which is an isomorphism for sufficiently fine covers. (A cover is *sufficiently fine* if $H^i(U_j, \mathcal{F}) = 0$ for all $i > 0$, for every j and for every sheaf \mathcal{F}). In the category of schemes, for example, any cover by open affine schemes has this property. What this means is that if one can find a finite fine enough cover of X , sheaf cohomology becomes computable by a finite process. In fact in [?], this is how the cohomology of projective space is explicitly calculated.

References

- [1] Grothendieck, A. *Sur quelques points d'algèbre homologique*, Tôhoku Math. J., Second Series, 9 (1957), 119–221.
- [2] Hartshorne, R. *Algebraic Geometry*, Springer-Verlag Graduate Texts in Mathematics 52, 1977