



Math for the people, by the people.

Hurwitz genus formula

Canonical name	HurwitzGenusFormula
Date of creation	2013-03-22 15:57:15
Last modified on	2013-03-22 15:57:15
Owner	alozano (2414)
Last modified by	alozano (2414)
Numerical id	6
Author	alozano (2414)
Entry type	Theorem
Classification	msc 14H99
Related topic	RiemannRochTheorem
Related topic	EllipticCurve

The following formula, due to Hurwitz, is extremely useful when trying to compute the genus of an algebraic curve. In this entry K is a perfect field (i.e. every algebraic extension of K is separable). Recall that a non-constant map of curves $\psi : C_1 \rightarrow C_2$ over K is separable if the extension of function fields $K(C_1)/\psi^*K(C_2)$ is a separable extension of fields.

Theorem (Hurwitz Genus Formula). *Let C_1 and C_2 be two smooth curves defined over K of genus g_1 and g_2 , respectively. Let $\psi : C_1 \rightarrow C_2$ be a non-constant and separable map. Then*

$$2g_1 - 2 \geq (\deg \psi)(2g_2 - 2) + \sum_{P \in C_1} (e_\psi(P) - 1)$$

where $e_\psi(P)$ is the ramification index of ψ at P . Moreover, there is equality if and only if either $\text{char}(K) = 0$ or $\text{char}(K) = p > 0$ and p does not divide $e_\psi(P)$ for all $P \in C_1$.

Example. As an application of the Hurwitz genus formula, we show that an elliptic curve $E : y^2 = x(x - \alpha)(x - \beta)$ defined over a field K of characteristic 0 has genus 1. Notice that the fact that E is an elliptic curve over K implies that $0, \alpha$ and β are distinct elements of K , otherwise E would be a singular curve. We define a map:

$$\psi : E \rightarrow \mathbb{P}^1, \quad [x, y, z] \mapsto [x, z]$$

and notice that $[0, 1, 0]$, the “point at infinity” of E , maps to $[1, 0]$, the point at infinity of \mathbb{P}^1 . The degree of this map is 2: generically every point in \mathbb{P}^1 has two preimages, namely $[x, y, z]$ and $[x, -y, z]$. Moreover, the genus of \mathbb{P}^1 is 0 and the map ψ is ramified exactly at 4 points, namely $P_1 = [0, 0, 1]$, $P_2 = [\alpha, 0, 1]$, $P_3 = [\beta, 0, 1]$ and the point at infinity. It is easily checked that the ramification index at each point is $e_\psi(P_i) = 2$. Hence, the Hurwitz formula reads:

$$2g_1 - 2 = 2(2 \cdot 0 - 2) + \sum_{i=1}^4 (e_\psi(P_i) - 1) = -4 + 4 = 0.$$

We conclude that $g_1 = 1$, as claimed.