



planetmath.org

Math for the people, by the people.

morphisms between bound quivers

Canonical name	MorphismsBetweenBoundQuivers
Date of creation	2013-03-22 19:17:07
Last modified on	2013-03-22 19:17:07
Owner	joking (16130)
Last modified by	joking (16130)
Numerical id	5
Author	joking (16130)
Entry type	Definition
Classification	msc 14L24

Let (Q, I) and (Q', I') be <http://planetmath.org/AdmissibleIdealsBoundQuiverAndItsAlgs> quivers over the same base field k .

Definition. A morphism $F : Q \rightarrow Q'$ is said to be **bounded by** (I, I') if <http://planetmath.org/MorphismsOfPathAlgebrasInducedFromMorphismsOfQuivers> the induced linear map $\overline{F} : kQ \rightarrow kQ'$ is such that

$$\overline{F}(I) \subseteq I'.$$

In this case we write

$$F : (Q, I) \rightarrow (Q', I')$$

and we say that F is a **morphism of bound quivers**.

If $F : (Q, I) \rightarrow (Q', I')$ is a morphism of bound quivers, then $\overline{F} : kQ \rightarrow kQ'$ induces a linear map

$$\overline{\overline{F}} : kQ/I \rightarrow kQ'/I'.$$

Furthermore, if F_0 is injective, then \overline{F} is a homomorphism of algebras (see <http://planetmath.org/MorphismsOfPathAlgebrasInducedFromMorphismsOfQuivers> this entry for details) and thus $\overline{\overline{F}}$ is a homomorphism of algebras.