



planetmath.org

Math for the people, by the people.

étale fundamental group

Canonical name	etaleFundamentalGroup
Date of creation	2013-03-22 14:11:46
Last modified on	2013-03-22 14:11:46
Owner	archibal (4430)
Last modified by	archibal (4430)
Numerical id	5
Author	archibal (4430)
Entry type	Definition
Classification	msc 14F35
Related topic	EtaleMorphism
Related topic	TopologicalSpace
Related topic	FundamentalGroup
Related topic	ClassificationOfCoveringSpaces

Recall that in topology, the fundamental group $\pi_1(T)$ of a connected topological space T is defined to be the group of loops based at a point modulo homotopy. When one wants to obtain something similar in the algebraic category, this definition encounters problems. One cannot simply attempt to use the same definition, since the result will be wrong if one is working in positive characteristic. More to the point, the topology on a scheme fails to capture much of the structure of the scheme. Simply choosing the “loop” to be an algebraic curve is not appropriate either, since in the most familiar case (over the complex numbers) such a “loop” has two real dimensions rather than one.

In the classification of covering spaces, it is shown that the fundamental group is exactly the group of deck transformations of the universal covering space. This is more promising: surjective étale morphisms are the appropriate generalization of covering spaces. Unfortunately, the universal covering space is often an infinite covering of the original space, which is unlikely to yield anything manageable in the algebraic category. Finite coverings, on the other hand are tractable, so one can define the algebraic fundamental group as an inverse limit of automorphism groups. Note also that finite étale morphisms are closed maps as well as open maps.

Definition 1 *Let X be a scheme, and let x be a geometric point of X . Then let C be the category of pairs (Y, π) such that $\pi: Y \rightarrow X$ is a finite étale morphism. Morphisms $(Y, \pi) \rightarrow (Y', \pi')$ in this category are morphisms $Y \rightarrow Y'$ as schemes over X . If Y' factors through Y as $Y' \rightarrow Y \rightarrow X$ then we obtain a morphism from $\text{Aut}_C(Y) \rightarrow \text{Aut}_C(Y')$. This allows us to construct the étale fundamental group*

$$\pi_1(X, x) = \varprojlim_{Y \in C} \text{Aut}_C(Y).$$

This explanation follows [?].

References

- [1] James Milne, *Lectures on Étale Cohomology*, <http://www.jmilne.org/math/CourseNotes/math732.html> online course notes.