



planetmath.org

Math for the people, by the people.

ideal

| | |
|------------------|---------------------|
| Canonical name | Ideal |
| Date of creation | 2013-03-22 11:49:27 |
| Last modified on | 2013-03-22 11:49:27 |
| Owner | djao (24) |
| Last modified by | djao (24) |
| Numerical id | 18 |
| Author | djao (24) |
| Entry type | Definition |
| Classification | msc 14K99 |
| Classification | msc 16D25 |
| Classification | msc 11N80 |
| Classification | msc 13A15 |
| Classification | msc 54C05 |
| Classification | msc 54C08 |
| Classification | msc 54J05 |
| Classification | msc 54D99 |
| Classification | msc 54E05 |
| Classification | msc 54E15 |
| Classification | msc 54E17 |
| Related topic | Subring |
| Related topic | PrimeIdeal |
| Defines | left ideal |
| Defines | right ideal |
| Defines | 2-sided ideal |
| Defines | two-sided ideal |

Let R be a ring. A *left ideal* (resp., *right ideal*) I of R is a nonempty subset $I \subset R$ such that:

- $a - b \in I$ for all $a, b \in I$
- $r \cdot a \in I$ (resp. $a \cdot r \in I$) for all $a \in I$ and $r \in R$

A *two-sided ideal* is a left ideal I which is also a right ideal. If R is a commutative ring, then these three notions of ideal are equivalent. Usually, the word “ideal” by itself means two-sided ideal.

The name “ideal” comes from the study of number theory. When the failure of unique factorization in number fields was first noticed, one of the solutions was to work with so-called “ideal numbers” in which unique factorization did hold. These “ideal numbers” were in fact ideals, and in Dedekind domains, unique factorization of ideals does indeed hold. The term “ideal number” is no longer used; the term “ideal” has replaced and generalized it.