



Math for the people, by the people.

## Picard group

Canonical name	PicardGroup
Date of creation	2013-03-22 12:52:30
Last modified on	2013-03-22 12:52:30
Owner	alozano (2414)
Last modified by	alozano (2414)
Numerical id	6
Author	alozano (2414)
Entry type	Definition
Classification	msc 14-00
Synonym	divisor class group

The *Picard group* of a variety, scheme, or more generally locally ringed space  $(X, O_X)$  is the group of locally free  $O_X$  modules of rank 1 with tensor product over  $O_X$  as the operation, usually denoted by  $\text{Pic}(X)$ . Alternatively, the Picard group is the group of isomorphism classes of invertible sheaves on  $X$ , under tensor products.

It is not difficult to see that  $\text{Pic}(X)$  is isomorphic to  $H^1(X, O_X^*)$ , the first sheaf cohomology group of the multiplicative sheaf  $O_X^*$  which consists of the units of  $O_X$ .

Finally, let  $\text{CaCl}(X)$  be the group of Cartier divisors on  $X$  modulo linear equivalence. If  $X$  is an integral scheme then the groups  $\text{CaCl}(X)$  and  $\text{Pic}(X)$  are isomorphic. Furthermore, if we let  $\text{Cl}(X)$  be the class group of Weil divisors (divisors modulo principal divisors) and  $X$  is a noetherian, integral and separated locally factorial scheme, then there is a natural isomorphism  $\text{Cl}(X) \cong \text{Pic}(X)$ . Thus, the Picard group is sometimes called the *divisor class group* of  $X$ .