



planetmath.org

Math for the people, by the people.

intersection divisor for a quartic

Canonical name	IntersectionDivisorForAQuartic
Date of creation	2013-03-22 15:44:59
Last modified on	2013-03-22 15:44:59
Owner	alozano (2414)
Last modified by	alozano (2414)
Numerical id	5
Author	alozano (2414)
Entry type	Definition
Classification	msc 14C20
Defines	hyperflex
Defines	flex

Let C be a non-singular curve in the plane, defined over an algebraically closed field K , and given by a polynomial $f(x, y) = 0$ of degree 4 (i.e. C is a quartic). Let L be a (rational) line in the plane K^2 . The intersection divisor of C and L is of the form:

$$(L \cdot C) = P_1 + P_2 + P_3 + P_4$$

where P_i , $i = 1, 2, 3, 4$, are points in $C(K)$. There are five possibilities:

1. The generic position: all the points P_i are distinct.
2. L is tangent to C : there exist indices $1 \leq i \neq j \leq 4$ such that $P_i = P_j$. Without loss of generality we may assume $P_1 = P_2$ and $(L \cdot C) = 2P_1 + P_3 + P_4$, and $P_3 \neq P_4$.
3. L is bitangent to C when $P_1 = P_2$ and $P_3 = P_4$ but $P_1 \neq P_3$. It may be shown that if $\text{char}(K) \neq 2$ then C has exactly 28 bitangent lines.
4. L intersects C at exactly two points, thus $P_1 = P_2 = P_3 \neq P_4$. The point P_1 is called a *flex*.
5. L intersects C at exactly one point and $P_1 = P_2 = P_3 = P_4$. This point is called a *hyperflex*. A quartic C may not have any hyperflex.

References

- [1] S. Flon, R. Oyono, C. Ritzenthaler, *Fast addition on non-hyperelliptic genus 3 curves*, can be found <http://eprint.iacr.org/2004/118.ps>.