



planetmath.org

Math for the people, by the people.

proof of determinant of the Vandermonde matrix

Canonical name	ProofOfDeterminantOfTheVandermondeMatrix
Date of creation	2013-03-22 15:44:50
Last modified on	2013-03-22 15:44:50
Owner	rspuzio (6075)
Last modified by	rspuzio (6075)
Numerical id	10
Author	rspuzio (6075)
Entry type	Proof
Classification	msc 15A57
Classification	msc 65F99
Classification	msc 65T50

To begin, note that the determinant of the $n \times n$ Vandermonde matrix (which we shall denote as ' Δ ') is a homogeneous polynomial of order $n(n-1)/2$ because every term in the determinant is, up to sign, the product of a zeroth power of some variable times the first power of some other variable, ..., the $n-1$ -st power of some variable and $0+1+\dots+(n-1) = n(n-1)/2$.

Next, note that if $a_i = a_j$ with $i \neq j$, then $\Delta = 0$ because two columns of the matrix would be equal. Since Δ is a polynomial, this implies that $a_i - a_j$ is a factor of Δ . Hence,

$$\Delta = C \prod_{1 \leq i < j \leq n} (a_j - a_i)$$

where C is some polynomial. However, since both Δ and the product on the right hand side have the same degree, C must have degree zero, i.e. C must be a constant. So all that remains is to determine the value of this constant.

One way to determine this constant is to look at the coefficient of the leading diagonal, $\prod_n (a_n)^{n-1}$. Since it equals 1 in both the determinant and the product, we conclude that $C = 1$, hence

$$\Delta = \prod_{1 \leq i < j \leq n} (a_j - a_i).$$