



planetmath.org

Math for the people, by the people.

cyclic decomposition theorem

Canonical name	CyclicDecompositionTheorem
Date of creation	2013-03-22 14:05:10
Last modified on	2013-03-22 14:05:10
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	16
Author	CWoo (3771)
Entry type	Theorem
Classification	msc 15A04
Synonym	T-admissible
Synonym	T -admissible
Related topic	CyclicSubspace
Defines	admissible subspace

Let k be a field, V a finite dimensional vector space over k and T a linear operator over V . Call a subspace $W \subseteq V$ *T-admissible* if W is T -invariant and for any polynomial $f(X) \in k[X]$ with $f(T)(v) \in W$ for $v \in V$, there is a $w \in W$ such that $f(T)(v) = f(T)(w)$.

Let W_0 be a proper T -admissible subspace of V . There are non zero vectors x_1, \dots, x_r in V with respective annihilator polynomials p_1, \dots, p_r such that

1. $V = W_0 \oplus Z(x_1, T) \oplus \dots \oplus Z(x_r, T)$ (See the cyclic subspace definition)
2. p_k divides p_{k-1} for every $k = 2, \dots, r$

Moreover, the integer r and the <http://planetmath.org/MinimalPolynomialEndomorphismminimalPolynomial> polynomials p_1, \dots, p_r are uniquely determined by (1),(2) and the fact that none of x_k is zero.

This is “one of the deepest results in linear algebra” (Hoffman & Kunze)