



direct sum of Hermitian and skew-Hermitian matrices

Canonical name	DirectSumOfHermitianAndSkewHermitianMatrices
Date of creation	2013-03-22 13:36:30
Last modified on	2013-03-22 13:36:30
Owner	mathcam (2727)
Last modified by	mathcam (2727)
Numerical id	5
Author	mathcam (2727)
Entry type	Example
Classification	msc 15A03
Classification	msc 15A57
Related topic	DirectSumOfEvenoddFunctionsExample

In this example, we show that any square matrix with complex entries can uniquely be decomposed into the sum of one Hermitian matrix and one skew-Hermitian matrix. A fancy way to say this is that complex square matrices is the direct sum of Hermitian and skew-Hermitian matrices.

Let us denote the vector space (over \mathbb{C}) of complex square $n \times n$ matrices by M . Further, we denote by M_+ respectively M_- the vector subspaces of Hermitian and skew-Hermitian matrices. We claim that

$$M = M_+ \oplus M_-. \quad (1)$$

Since M_+ and M_- are vector subspaces of M , it is clear that $M_+ + M_-$ is a vector subspace of M . Conversely, suppose $A \in M$. We can then define

$$\begin{aligned} A_+ &= \frac{1}{2}(A + A^*), \\ A_- &= \frac{1}{2}(A - A^*). \end{aligned}$$

Here $A^* = \overline{A}^T$, and \overline{A} is the complex conjugate of A , and A^T is the transpose of A . It follows that A_+ is Hermitian and A_- is anti-Hermitian. Since $A = A_+ + A_-$, any element in M can be written as the sum of one element in M_+ and one element in M_- . Let us check that this decomposition is unique. If $A \in M_+ \cap M_-$, then $A = A^* = -A$, so $A = 0$. We have established equation ??.

Special cases

- In the special case of 1×1 matrices, we obtain the decomposition of a complex number into its real and imaginary components.
- In the special case of real matrices, we obtain the decomposition of a $n \times n$ matrix into a symmetric matrix and anti-symmetric matrix.