



planetmath.org

Math for the people, by the people.

matrix unit

| | |
|------------------|--------------------------|
| Canonical name | MatrixUnit |
| Date of creation | 2013-03-22 18:30:35 |
| Last modified on | 2013-03-22 18:30:35 |
| Owner | CWoo (3771) |
| Last modified by | CWoo (3771) |
| Numerical id | 8 |
| Author | CWoo (3771) |
| Entry type | Definition |
| Classification | msc 15A30 |
| Classification | msc 16S50 |
| Related topic | ElementaryMatrix |
| Defines | full set of matrix units |

A *matrix unit* is a matrix (over some ring with 1) whose entries are all 0 except in one cell, where it is 1.

For example, among the 3×2 matrices,

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{pmatrix}$$

are the matrix units.

Let A and B be $m \times n$ and $p \times q$ matrices over R , and U_{ij} an $n \times p$ matrix unit (over R). Then

1. AU_{ij} is the $m \times p$ matrix whose j th column is the i th column of A , and 0 everywhere else, and
2. $U_{ij}B$ is the $n \times q$ matrix whose i th row is the j th row of B and 0 everywhere else.

Remarks. Let $M = M_{m \times n}(R)$ be the set of all m by n matrices with entries in a ring R (with 1). Denote U_{ij} the matrix unit in M whose cell (i, j) is 1.

- M is a (left or right) R -module generated by the $m \times n$ matrix units.
- When $m = n$, M has the structure of an algebra over R . The matrix units have the following properties:

1. $U_{ij}U_{k\ell} = \delta_{jk}U_{i\ell}$, and
2. $U_{11} + \cdots + U_{nn} = I_n$,

where δ_{ij} is the Kronecker delta and I_n is the identity matrix. Note that the U_{ii} form a complete set of pairwise orthogonal idempotents, meaning $U_{ii}U_{ii} = U_{ii}$ and $U_{ii}U_{jj} = 0$ if $i \neq j$.

- In general, in a matrix ring S (consisting of, say, all $n \times n$ matrices), any set of n matrices satisfying the two properties above is called a *full set of matrix units* of S .
- For example, if $\{U_{ij} \mid 1 \leq i, j \leq 2\}$ is the set of 2×2 matrix units over \mathbb{R} , then for any invertible matrix T , $\{TU_{ij}T^{-1} \mid 1 \leq i, j \leq 2\}$ is a full set of matrix units.

- If we embed R as a subring of $M_n(R)$, then R is the centralizer of the matrix units of $M_n(R)$, meaning that the only elements in $M_n(R)$ that commute with the matrix units are the elements in R .

References

- [1] T. Y. Lam, *Lectures on Modules and Rings*, Springer, New York, 1998.