



planetmath.org

Math for the people, by the people.

block determinants

|                  |   |
|------------------|---|
| Canonical name   | BlockDeterminants                       |
| Date of creation | 2013-03-22 15:25:57                     |
| Last modified on | 2013-03-22 15:25:57                     |
| Owner            | georgiosl (7242)                        |
| Last modified by | georgiosl (7242)                        |
| Numerical id     | 20                                      |
| Author           | georgiosl (7242)                        |
| Entry type       | Theorem                                 |
| Classification   | msc 15A15                               |
| Related topic    | SchurComplement                         |
| Related topic    | DeterminantsOfSomeMatricesOfSpecialForm |

If  $A$  and  $D$  are square matrices

- If  $A^{-1}$  exists, then

$$\det \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \det(A) \det(D - CA^{-1}B)$$

- If  $D^{-1}$  exists, then

$$\det \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \det(D) \det(A - BD^{-1}C)$$

The matrices  $D - CA^{-1}B$  and  $A - BD^{-1}C$  are called the Schur complements of  $A$  and  $D$ , respectively.

Mention that

- If  $A, D$  are square matrices, then

$$\det \begin{pmatrix} A & B \\ O & D \end{pmatrix} = \det(A) \det(D)$$

, where  $O$  is a zero matrix.

- Also we have that

$$\det \begin{pmatrix} A & O \\ O & B \end{pmatrix} = \det(A) \det(B).$$

- Another useful result for block determinants is the following.

As  $J = \begin{pmatrix} O & I \\ -I & O \end{pmatrix}$  is a symplectic matrix, we have that  $\det J = 1$ .

Using now the fact that  $\det MN = \det(M) \det(N)$  for any  $M, N$  square matrices, we have that

$$\det \begin{pmatrix} O & A \\ B & O \end{pmatrix} = \det \begin{pmatrix} O & A \\ B & O \end{pmatrix} \det J = -\det(A) \det(B)$$

This holds for any square matrices  $A, B$  and for the last point  $A, B$  have also the same order. They do not need to be invertible.