



Math for the people, by the people.

trace

| | |
|------------------|---------------------|
| Canonical name | Trace |
| Date of creation | 2013-03-22 12:17:57 |
| Last modified on | 2013-03-22 12:17:57 |
| Owner | mhale (572) |
| Last modified by | mhale (572) |
| Numerical id | 10 |
| Author | mhale (572) |
| Entry type | Definition |
| Classification | msc 15A15 |
| Classification | msc 15A04 |
| Related topic | FrobeniusMatrixNorm |

The *trace* $\text{Tr}(A)$ of a square matrix A is defined to be the sum of the diagonal entries of A . It satisfies the following formulas:

- $\text{Tr}(A + B) = \text{Tr}(A) + \text{Tr}(B)$
- $\text{Tr}(AB) = \text{Tr}(BA)$ ()

where A and B are square matrices of the same size.

The *trace* $\text{Tr}(T)$ of a linear transformation $T: V \longrightarrow V$ from any finite dimensional vector space V to itself is defined to be the trace of any matrix representation of T with respect to a basis of V . This scalar is independent of the choice of basis of V , and in fact is equal to the sum of the eigenvalues of T (over a splitting field of the characteristic polynomial), including multiplicities.

The following link presents some examples for calculating the trace of a matrix.

A *trace* on a C^* -algebra A is a positive linear functional $\phi: A \rightarrow \mathbb{C}$ that has the .