



planetmath.org

Math for the people, by the people.

Jordan canonical form theorem

Canonical name	JordanCanonicalFormTheorem
Date of creation	2013-03-22 12:59:21
Last modified on	2013-03-22 12:59:21
Owner	Mathprof (13753)
Last modified by	Mathprof (13753)
Numerical id	16
Author	Mathprof (13753)
Entry type	Theorem
Classification	msc 15A18
Synonym	Jordan canonical form
Related topic	PartitionedMatrix
Related topic	SimultaneousUpperTriangularBlockDiagonalizationOfCommutingMatrices
Related topic	Diagonalizable2
Defines	Jordan block
Defines	Jordan matrix

A **Jordan block** or **Jordan matrix** is a matrix of the form

$$\begin{pmatrix} \lambda & 1 & 0 & \cdots & 0 \\ 0 & \lambda & 1 & \cdots & 0 \\ 0 & 0 & \lambda & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda \end{pmatrix}$$

with a constant value λ along the diagonal and 1's on the superdiagonal. Some texts use the 1's on the subdiagonal instead.

Theorem. *Let V be a finite-dimensional vector space over a field F and $t : V \rightarrow V$ be a linear transformation. Then, if the characteristic polynomial factors completely over F , there will exist a basis of V with respect to which the matrix of t is of the form*

$$\begin{pmatrix} J_1 & 0 & \cdots & 0 \\ 0 & J_2 & \cdots & 0 \\ & & \cdots & \\ 0 & 0 & \cdots & J_k \end{pmatrix}$$

where each J_i is a Jordan block in which $\lambda = \lambda_i$.

The matrix in Theorem 1 is called a *Jordan canonical form* for the transformation t .