



planetmath.org

Math for the people, by the people.

Frobenius method

Canonical name	FrobeniusMethod
Date of creation	2013-03-22 17:43:49
Last modified on	2013-03-22 17:43:49
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	18
Author	pahio (2872)
Entry type	Topic
Classification	msc 15A06
Classification	msc 34A05
Synonym	method of Frobenius
Related topic	FuchsianSingularity
Related topic	BesselsEquation
Related topic	SpecialCasesOfHypergeometricFunction
Defines	indicial equation

Let us consider the linear homogeneous differential equation

$$\sum_{\nu=0}^n k_{\nu}(x)y^{(n-\nu)}(x) = 0$$

of <http://planetmath.org/DifferentialEquationorder> n . If the coefficient functions $k_{\nu}(x)$ are continuous and the coefficient $k_0(x)$ of the highest <http://planetmath.org/HigherOrderDerivativesorder> derivative does not vanish on a certain interval (resp. a <http://planetmath.org/Domain2domain> in \mathbb{C}), then all solutions $y(x)$ are continuous on this interval (resp.). If all coefficients have the continuous derivatives up to a certain , the same concerns the solutions.

If, instead, $k_0(x)$ vanishes in a point x_0 , this point is in general a singular point. After dividing the differential equation by $k_0(x)$ and then getting the form

$$y^{(n)}(x) + \sum_{\nu=1}^n c_{\nu}(x)y^{(n-\nu)}(x) = 0,$$

some new coefficients $c_{\nu}(x)$ are discontinuous in the singular point. However, if the discontinuity is so, that the products

$$(x - x_0)c_1(x), \quad (x - x_0)^2c_2(x), \quad \dots, \quad (x - x_0)^nc_n(x)$$

are continuous, and analytic in x_0 , the point x_0 is a regular singular point of the differential equation.

We introduce the so-called *Frobenius method* for finding solution functions in a neighbourhood of the regular singular point x_0 , confining us to the case of a <http://planetmath.org/DifferentialEquationsecond> order differential equation. When we use the <http://planetmath.org/Divisionquotient> forms

$$(x - x_0)c_1(x) := \frac{p(x)}{r(x)}, \quad (x - x_0)^2c_2(x) := \frac{q(x)}{r(x)},$$

where $r(x)$, $p(x)$ and $q(x)$ are analytic in a neighbourhood of x_0 and $r(x) \neq 0$, our differential equation reads

$$(x - x_0)^2r(x)y''(x) + (x - x_0)p(x)y'(x) + q(x)y(x) = 0. \quad (1)$$

Since a change $x - x_0 \mapsto x$ of variable brings to the case that the singular point is the origin, we may suppose such a starting situation. Thus we can study the equation

$$x^2 r(x) y''(x) + x p(x) y'(x) + q(x) y(x) = 0, \quad (2)$$

where the coefficients have the converging power series expansions

$$r(x) = \sum_{n=0}^{\infty} r_n x^n, \quad p(x) = \sum_{n=0}^{\infty} p_n x^n, \quad q(x) = \sum_{n=0}^{\infty} q_n x^n \quad (3)$$

and

$$r_0 \neq 0.$$

In the Frobenius method one examines whether the equation (2) allows a series solution of the form

$$y(x) = x^s \sum_{n=0}^{\infty} a_n x^n = a_0 x^s + a_1 x^{s+1} + a_2 x^{s+2} + \dots, \quad (4)$$

where s is a constant and $a_0 \neq 0$.

Substituting (3) and (4) to the differential equation (2) converts the left hand to

$$\begin{aligned} & [r_0 s(s-1) + p_0 s + q_0] a_0 x^s + \\ & [[r_0(s+1)s + p_0(s+1) + q_0] a_1 + [r_1 s(s-1) + p_1 s + q_1] a_0] x^{s+1} + \\ & [[r_0(s+2)(s+1) + p_0(s+2) + q_0] a_2 + [r_1(s+1)s + p_1(s+1) + q_1] a_1 + [r_2 s(s-1) + p_2 s + q_2] a_0] x^{s+2} + \dots \end{aligned}$$

Our equation seems clearer when using the notations $f_\nu(s) := r_\nu s(s-1) + p_\nu s + q_\nu$:

$$f_0(s) a_0 x^s + [f_0(s+1) a_1 + f_1(s) a_0] x^{s+1} + [f_0(s+2) a_2 + f_1(s+1) a_1 + f_2(s) a_0] x^{s+2} + \dots = 0 \quad (5)$$

Thus the condition of satisfying the differential equation by (4) is the infinite system of equations

$$\begin{cases} f_0(s) a_0 = 0 \\ f_0(s+1) a_1 + f_1(s) a_0 = 0 \\ f_0(s+2) a_2 + f_1(s+1) a_1 + f_2(s) a_0 = 0 \\ \dots \quad \dots \quad \dots \end{cases} \quad (6)$$

In the first , since $a_0 \neq 0$, the *indicial equation*

$$f_0(s) \equiv r_0 s^2 + (p_0 - r_0)s + q_0 = 0 \quad (7)$$

must be satisfied. Because $r_0 \neq 0$, this quadratic equation determines for s two values, which in special case may coincide.

The first of the equations (6) leaves $a_0 (\neq 0)$ arbitrary. The next linear equations in a_n allow to solve successively the constants a_1, a_2, \dots provided that the first coefficients $f_0(s+1), f_0(s+2), \dots$ do not vanish; this is evidently the case when the <http://planetmath.org/Equationroots> of the indicial equation don't differ by an integer (e.g. when the are complex conjugates or when s is the having greater real part). In any case, one obtains at least for one of the of the indicial equation the definite values of the coefficients a_n in the series (4). It is not hard to show that then this series converges in a neighbourhood of the origin.

For obtaining the solution of the differential equation (2) it suffices to have only one solution $y_1(x)$ of the form (4), because another solution $y_2(x)$, linearly independent on $y_1(x)$, is gotten via mere integrations; then it is possible in the cases $s_1 - s_2 \in \mathbb{Z}$ that $y_2(x)$ has no expansion of the form (4).

References

- [1] PENTTI LAASONEN: *Matemaattisia erikoisfunktioita*. Handout No. 261. Teknillisen Korkeakoulun Ylioppilaskunta; Otaniemi, Finland (1969).