



planetmath.org

Math for the people, by the people.

dimension formulae for vector spaces

Canonical name	DimensionFormulaeForVectorSpaces
Date of creation	2013-03-22 16:31:19
Last modified on	2013-03-22 16:31:19
Owner	yark (2760)
Last modified by	yark (2760)
Numerical id	25
Author	yark (2760)
Entry type	Feature
Classification	msc 15A03

In this entry we look at various formulae involving the dimension of a vector space.

Throughout this entry, K will be a field, and V and W will be vector spaces over K . The dimension of a vector space U over K will be denoted by $\dim(U)$, or by $\dim_K(U)$ if the ground field needs to be emphasized.

All of these formulae potentially involve infinite cardinals, so the reader should have a basic knowledge of cardinal arithmetic in order to understand them in full generality.

Subspaces

If S and T are subspaces of V , then

$$\dim(S) + \dim(T) = \dim(S \cap T) + \dim(S + T).$$

Rank-nullity theorem

The rank-nullity theorem states that if $\phi: V \rightarrow W$ is a linear mapping, then the dimension of V is the sum of the dimensions of the image and kernel of ϕ :

$$\dim(V) = \dim(\text{Im } \phi) + \dim(\text{Ker } \phi).$$

In particular, if U is a subspace of V then

$$\dim(V) = \dim(V/U) + \dim(U).$$

The rank-nullity theorem can also be stated in terms of short exact sequences: if

$$0 \longrightarrow U \longrightarrow V \longrightarrow W \longrightarrow 0$$

is a short exact sequence of vector spaces over K , then

$$\dim(V) = \dim(U) + \dim(W).$$

This can be generalized to infinite exact sequences: if

$$\cdots \longrightarrow V_{n+1} \longrightarrow V_n \longrightarrow V_{n-1} \longrightarrow \cdots$$

is an exact sequence of vector spaces over K , then

$$\sum_{n \text{ even}} \dim(V_n) = \sum_{n \text{ odd}} \dim(V_n).$$

(This is indeed a generalization, because any finite exact sequence of vector spaces terminating with 0 at both ends can be considered as an infinite exact sequence in which all remaining terms are 0.)

Direct sums

If $(V_i)_{i \in I}$ is a family of vector spaces over K , then

$$\dim\left(\bigoplus_{i \in I} V_i\right) = \sum_{i \in I} \dim(V_i).$$

Cardinality of a vector space

The cardinality of a vector space is determined by its dimension and the cardinality of the ground field:

$$|V| = \begin{cases} |K|^{\dim(V)}, & \text{if } \dim(V) \text{ is finite;} \\ \max\{|K|, \dim(V)\}, & \text{if } \dim(V) \text{ is infinite.} \end{cases}$$

The effect of the above formula is somewhat different depending on whether K is <http://planetmath.org/FiniteField> finite or infinite. If K is finite, then it reduces to

$$|V| = \begin{cases} |K|^{\dim(V)}, & \text{if } \dim(V) \text{ is finite;} \\ \dim(V), & \text{if } \dim(V) \text{ is infinite.} \end{cases}$$

If K is infinite, then it can be expressed as

$$|V| = \begin{cases} 1, & \text{if } \dim(V) = 0; \\ |K|, & \text{if } 0 < \dim(V) \leq |K|; \\ \dim(V), & \text{if } \dim(V) \geq |K|. \end{cases}$$

Change of ground field

If F is a subfield of K , then V can be considered as a vector space over F . The dimensions of V over K and F are related by the formula

$$\dim_F(V) = [K : F] \cdot \dim_K(V).$$

In this formula, $[K : F]$ is the degree of the field extension K/F , that is, the dimension of K considered as a vector space over F .

Space of functions into a vector space

If S is any set, then the set K^S of all functions from S into K becomes a vector space over K if we define the operations pointwise, that is, $(f+g)(x) = f(x) + g(x)$ and $(\lambda f)(x) = \lambda f(x)$ for all $f, g \in K^S$, all $x \in S$, and all $\lambda \in K$. The dimension of this vector space is given by

$$\dim(K^S) = \begin{cases} |S|, & \text{if } S \text{ is finite;} \\ |K|^{|S|}, & \text{if } S \text{ is infinite.} \end{cases}$$

The case where S is infinite is not straightforward to prove. Proofs can be found in books by Baer[?] and Jacobson[?], among others.

More generally, we can consider the space V^S , which is really just the <http://planetmath.org/DirectProduct> direct product of copies of V indexed by S . We get

$$\dim(V^S) = \begin{cases} 0, & \text{if } \dim(V) = 0; \\ |S| \cdot \dim(V), & \text{if } S \text{ is finite;} \\ |V|^{|S|}, & \text{otherwise.} \end{cases}$$

Dual space

Given any basis B of V , the dual space V^* is isomorphic to K^B via the mapping $f \mapsto f|_B$. So the formula of the previous section can be applied to give a formula for the dimension of V^* :

$$\dim(V^*) = \begin{cases} \dim(V), & \text{if } \dim(V) \text{ is finite;} \\ |K|^{\dim(V)}, & \text{if } \dim(V) \text{ is infinite.} \end{cases}$$

In particular, this formula implies that V is isomorphic to V^* if and only if V is finite-dimensional. (Students who are familiar with the fact that an infinite-dimensional Banach space can be isomorphic to its dual are sometimes surprised to learn that an infinite-dimensional vector space cannot be isomorphic to its dual, for a Banach space is surely a vector space. But the

term *dual* is used in different senses in these two statements, so there is no contradiction. In the theory of Banach spaces one is usually only interested in the *continuous* linear functionals, and the resulting ‘continuous’ dual is a subspace of the full dual used in the above formula.)

Space of linear mappings

The set $\text{Hom}_K(V, W)$ of all linear mappings from V into W is itself a vector space over K , with the operations defined in the obvious way, namely $(f + g)(x) = f(x) + g(x)$ and $(\lambda f)(x) = \lambda f(x)$ for all $f, g \in \text{Hom}_K(V, W)$, all $x \in V$, and all $\lambda \in K$. The dual space $V^* = \text{Hom}_K(V, K)$ considered in the previous section is a special case of this. For any basis B of V , the mapping $f \mapsto f|_B$ defines an isomorphism between $\text{Hom}_K(V, W)$ and W^B , so that from an earlier section we get

$$\dim(\text{Hom}_K(V, W)) = \begin{cases} 0, & \text{if } \dim(W) = 0; \\ \dim(V) \cdot \dim(W), & \text{if } \dim(V) \text{ is finite;} \\ |W|^{\dim(V)}, & \text{otherwise.} \end{cases}$$

In the special case $W = V$ this can be simplified to

$$\dim(\text{End}_K(V)) = \begin{cases} \dim(V)^2, & \text{if } \dim(V) \text{ is finite;} \\ |K|^{\dim(V)}, & \text{otherwise.} \end{cases}$$

Tensor products

The dimension of the <http://planetmath.org/TensorProduct> tensor product of V and W is given by

$$\dim(V \otimes W) = \dim(V) \cdot \dim(W).$$

Banach spaces

The dimension of a Banach space, considered as a vector space, is sometimes called the *Hamel dimension*, in order to distinguish it from other concepts of dimension. For an infinite-dimensional Banach space B we have

$$\dim(B) = |B|.$$

The tricky part of establishing this formula is to show that the dimension is always at least the cardinality of the continuum. A short proof of this is given in a paper by Lacey[?].

The above formula suggests that Hamel dimension is not a very useful concept for infinite-dimensional Banach spaces, which is indeed the case. Nonetheless, it is interesting to see how Hamel dimension relates to the usual concept of dimension in Hilbert spaces. If H is a Hilbert space, and d is its dimension (meaning the cardinality of an orthonormal basis), then the Hamel dimension $\dim(H)$ is given by

$$\dim(H) = \begin{cases} d, & \text{if } d \text{ is finite;} \\ d^{\aleph_0}, & \text{if } d \text{ is infinite.} \end{cases}$$

References

- [1] Reinhold Baer, *Linear Algebra and Projective Geometry*, Academic Press, 1952.
- [2] Nathan Jacobson, *Lectures in Abstract Algebra*, Volume II: *Linear Algebra*, D. Van Nostrand Company Inc., 1953.
- [3] H. Elton Lacey, *The Hamel Dimension of any Infinite Dimensional Separable Banach Space is c* , Amer. Math. Mon. 80 (1973), 298.