



planetmath.org

Math for the people, by the people.

Ostrowski theorem

Canonical name	OstrowskiTheorem
Date of creation	2013-03-22 15:36:29
Last modified on	2013-03-22 15:36:29
Owner	Andrea Ambrosio (7332)
Last modified by	Andrea Ambrosio (7332)
Numerical id	22
Author	Andrea Ambrosio (7332)
Entry type	Theorem
Classification	msc 15A42

Let A be a complex $n \times n$ matrix, $R_i = \sum_{j \neq i} |a_{ij}|$, $C_j = \sum_{i \neq j} |a_{ij}|$ $1 \leq i \leq n, 1 \leq j \leq n$. Let's consider, for any $\alpha \in (0, 1)$, the circles of this kind: $O_i = \{z \in \mathbf{C} : |z - a_{ii}| \leq R_i^\alpha C_i^{1-\alpha}\}$ $1 \leq i \leq n$.

Theorem (A. Ostrowski): For any $\alpha \in (0, 1)$, all the eigenvalues of A lie in the union of these n circles: $\sigma(A) \subseteq \bigcup_i O_i$.

Proof. If $R_i = 0$, the theorem says a_{ii} is an eigenvalue, which is obviously true. Let's then concentrate on the $R_i \neq 0$. By eigenvalue definition, we have:

$$(\lambda - a_{ii})x_i = \sum_{j \neq i} a_{ij}x_j$$

so that, recalling Hölder's inequality with $p = 1/\alpha$ and $q = 1/(1-\alpha)$ (to have $p, q > 1$, we must have $\alpha \in (0, 1)$)

$$\begin{aligned} |\lambda - a_{ii}| |x_i| &\leq \sum_{j \neq i} |a_{ij}| |x_j| \\ &= \sum_{j \neq i} |a_{ij}|^\alpha |a_{ij}|^{1-\alpha} |x_j| \\ &\leq \left(\sum_{j \neq i} (|a_{ij}|^\alpha)^{1/\alpha} \right)^\alpha \left(\sum_{j \neq i} (|a_{ij}|^{1-\alpha} |x_j|)^{1/(1-\alpha)} \right)^{1-\alpha} \\ &= \left(\sum_{j \neq i} |a_{ij}| \right)^\alpha \left(\sum_{j \neq i} |a_{ij}| |x_j|^{1/(1-\alpha)} \right)^{1-\alpha} \\ &= R_i^\alpha \left(\sum_{j \neq i} |a_{ij}| |x_j|^{1/(1-\alpha)} \right)^{1-\alpha} \end{aligned}$$

which means

$$\frac{|\lambda - a_{ii}|^{1/(1-\alpha)}}{R_i^{\alpha/(1-\alpha)}} |x_i|^{1/(1-\alpha)} \leq \sum_{j \neq i} |a_{ij}| |x_j|^{1/(1-\alpha)}$$

Summing over all i , one obtains

$$\sum_{i=1}^n \frac{|\lambda - a_{ii}|^{1/(1-\alpha)}}{R_i^{\alpha/(1-\alpha)}} |x_i|^{1/(1-\alpha)} \leq \sum_{i=1}^n \sum_{j \neq i} |a_{ij}| |x_j|^{1/(1-\alpha)} = \sum_{j=1}^n C_j |x_j|^{1/(1-\alpha)}$$

If, for each i , the coefficient of $|x_i|^{1/(1-\alpha)}$ in the first sum would be greater than the coefficient of the same term in the right-hand side, inequality couldn't hold. So we can conclude that at least one index p exists such as

$$\frac{|\lambda - a_{pp}|^{1/(1-\alpha)}}{R_p^{\alpha/(1-\alpha)}} \leq C_p$$

that is

$$|\lambda - a_{pp}| \leq R_p^\alpha C_p^{1-\alpha}$$

which is the thesis. □

Remarks:

The Gershgorin theorem is obtained as a limit for $\alpha \rightarrow 0$ or for $\alpha \rightarrow 1$; in other words, Ostrowski's theorem represents a kind of "continuous deformation" between the two Gershgorin rows and columns sets.

References

- [1] R. A. Horn, C. R. Johnson, *Matrix Analysis*, Cambridge University Press, 1985