



Math for the people, by the people.

## Gerstenhaber - Serezhkin theorem

Canonical name	GerstenhaberSerezhkinTheorem
Date of creation	2013-03-22 19:20:05
Last modified on	2013-03-22 19:20:05
Owner	kammerer (26336)
Last modified by	kammerer (26336)
Numerical id	7
Author	kammerer (26336)
Entry type	Theorem
Classification	msc 15A30
Related topic	BottaPierceWatkinsTheorem

Let  $\mathbb{F}$  be an arbitrary field. Consider  $\mathcal{M}_n(\mathbb{F})$ , the vector space of all  $n \times n$  matrices over  $\mathbb{F}$ . Define

- $\mathcal{N} = \{A \in \mathcal{M}_n(\mathbb{F}) : A \text{ is nilpotent}\},$
- $\mathcal{GL}_n(\mathbb{F}) = \{A \in \mathcal{M}_n(\mathbb{F}) : \det(A) \neq 0\},$
- $\mathcal{T} = \{A \in \mathcal{M}_n(\mathbb{F}) : A \text{ is strictly upper triangular}\}.$

Notice that  $\mathcal{T}$  is a linear subspace of  $\mathcal{M}_n(\mathbb{F})$ . Moreover,  $\mathcal{T} \subseteq \mathcal{N}$  and  $\dim \mathcal{T} = n(n-1)/2$ .

The Gerstenhaber – Serezhkin theorem on linear subspaces contained in the nilpotent cone [?, ?] reads as follows.

**Theorem 1** *Let  $\mathcal{L}$  be a linear subspace of  $\mathcal{M}_n(\mathbb{F})$ . Assume that  $\mathcal{L} \subseteq \mathcal{N}$ . Then*

- (i)  $\dim \mathcal{L} \leq n(n-1)/2,$
- (ii)  $\dim \mathcal{L} = n(n-1)/2$  if and only if there exists  $U \in \mathcal{GL}_n(\mathbb{F})$  such that  $\{UAU^{-1} : A \in \mathcal{L}\} = \mathcal{T}$ .

An alternative simple proof of inequality (i) can be found in [?].

## References

- [G] M. Gerstenhaber, On nilalgebras and linear varieties of nilpotent matrices, I, *Amer. J. Math.* **80**: 614–622 (1958).
- [M] B. Mathes, M. Omladič, H. Radjavi, Linear Spaces of Nilpotent Matrices, *Linear Algebra Appl.* **149**: 215–225 (1991).
- [S] V. N. Serezhkin, On linear transformations preserving nilpotency, *Vestsi Akad. Navuk BSSR Ser. Fiz.-Mat. Navuk* **1985**, no. 6: 46–50 (Russian).