



planetmath.org

Math for the people, by the people.

position vector

Canonical name	PositionVector
Date of creation	2013-03-22 15:25:15
Last modified on	2013-03-22 15:25:15
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	18
Author	pahio (2872)
Entry type	Definition
Classification	msc 15A72
Synonym	radius vector
Related topic	ExampleOfCurvatureSpaceCurve
Related topic	DyadProduct
Related topic	TiltCurve

In the space \mathbb{R}^3 , the vector

$$\vec{r} := (x, y, z) = x\vec{i} + y\vec{j} + z\vec{k}$$

directed from the origin to a point (x, y, z) is the *position vector* of this point. When the point is , \vec{r} a vector field and its

$$r := \sqrt{x^2 + y^2 + z^2}$$

a scalar .

The

- $\nabla \cdot \vec{r} = 3$
- $\nabla \times \vec{r} = \vec{0}$
- $\nabla r = \frac{\vec{r}}{r} = \vec{r}^0$
- $\nabla \frac{1}{r} = -\frac{\vec{r}}{r^3} = -\frac{\vec{r}^0}{r^2}$
- $\nabla^2 \frac{1}{r} = 0$

are valid, where \vec{r}^0 is the unit vector having the direction of \vec{r} .

If \vec{c} is a vector, $\vec{U}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ a vector function and $f: \mathbb{R} \rightarrow \mathbb{R}$ is a twice differentiable function, then the formulae

- $\nabla(\vec{c} \cdot \vec{r}) = \vec{c}$
- $\nabla \cdot (\vec{c} \times \vec{r}) = 0$
- $(\vec{U} \cdot \nabla) \vec{r} = \vec{U}$
- $(\vec{U} \times \nabla) \cdot \vec{r} = 0$
- $(\vec{U} \times \nabla) \times \vec{r} = -2\vec{U}$
- $\nabla f(r) = f'(r) \vec{r}^0$
- $\nabla^2 f(r) = f''(r) + \frac{2}{r} f'(r)$

hold.

References

- [1] K. VÄISÄLÄ: *Vektorianalyysi*. Werner Söderström Osakeyhtiö, Helsinki (1961).