



Math for the people, by the people.

proof of Bauer-Fike theorem

Canonical name	ProofOfBauerFikeTheorem
Date of creation	2013-03-22 15:33:08
Last modified on	2013-03-22 15:33:08
Owner	Andrea Ambrosio (7332)
Last modified by	Andrea Ambrosio (7332)
Numerical id	9
Author	Andrea Ambrosio (7332)
Entry type	Proof
Classification	msc 15A42

We can assume $\tilde{\lambda} \notin \sigma(A)$ (otherwise, we can choose $\lambda = \tilde{\lambda}$ and theorem is proven, since $\kappa_p(X) > 1$). Then $(A - \tilde{\lambda}I)^{-1}$ exists, so we can write:

$$\tilde{u} = (A - \tilde{\lambda}I)^{-1}r = X(D - \tilde{\lambda}I)^{-1}X^{-1}r$$

since A is diagonalizable; taking the <http://planetmath.org/VectorPnormp>-norm of both sides, we obtain:

$$\begin{aligned} 1 &= \|\tilde{u}\|_p \\ &= \|X(D - \tilde{\lambda}I)^{-1}X^{-1}r\|_p \leq \|X\|_p \|(D - \tilde{\lambda}I)^{-1}\|_p \|X^{-1}\|_p \|r\|_p \\ &= \kappa_p(X) \|(D - \tilde{\lambda}I)^{-1}\|_p \|r\|_p. \end{aligned}$$

But, since $(D - \tilde{\lambda}I)^{-1}$ is a diagonal matrix, the p-norm is easily computed, and yields:

$$\|(D - \tilde{\lambda}I)^{-1}\|_p = \max_{\|x\|_p \neq 0} \frac{\|(D - \tilde{\lambda}I)^{-1}x\|_p}{\|x\|_p} = \max_{\lambda \in \sigma(A)} \frac{1}{|\lambda - \tilde{\lambda}|} = \frac{1}{\min_{\lambda \in \sigma(A)} |\lambda - \tilde{\lambda}|}$$

whence:

$$\min_{\lambda \in \sigma(A)} |\lambda - \tilde{\lambda}| \leq \kappa_p(X) \|r\|_p.$$