



Math for the people, by the people.

## proof of block determinants

Canonical name	ProofOfBlockDeterminants
Date of creation	2013-03-22 15:27:49
Last modified on	2013-03-22 15:27:49
Owner	georgiosl (7242)
Last modified by	georgiosl (7242)
Numerical id	5
Author	georgiosl (7242)
Entry type	Proof
Classification	msc 15A15

If  $A^{-1}$  exists, then

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} I & O \\ CA^{-1} & I \end{pmatrix} \begin{pmatrix} A & B \\ O & D - CA^{-1}B \end{pmatrix}.$$

So

$$\det \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \det \begin{pmatrix} I & O \\ CA^{-1} & I \end{pmatrix} \det \begin{pmatrix} A & B \\ O & D - CA^{-1}B \end{pmatrix}.$$

Each of the first matrices in the decompositions are triangular. Hence their determinants equal 1. This means that the determinant of the original matrix equals the determinant of either of the second matrices in the decomposition. Therefore

$$\det \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \det(A) \det(D - CA^{-1}B).$$

The second formula follows by using a similar trick.