



Math for the people, by the people.

derivative of homogeneous function

Canonical name	DerivativeOfHomogeneousFunction
Date of creation	2013-03-22 14:45:05
Last modified on	2013-03-22 14:45:05
Owner	matte (1858)
Last modified by	matte (1858)
Numerical id	9
Author	matte (1858)
Entry type	Theorem
Classification	msc 15-00

Theorem 1. *Suppose $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a differentiable positively homogeneous function of degree r . Then $\frac{\partial f}{\partial x^i}$ is a positively homogeneous function of degree $r - 1$.*

Proof. By considering component functions if necessary, we can assume that $m = 1$. For $\lambda \in \mathbb{R}$, let M_λ be the multiplication map,

$$\begin{aligned} M_\lambda: \mathbb{R}^n &\rightarrow \mathbb{R}^n \\ v &\mapsto \lambda v. \end{aligned}$$

For $\lambda > 0$ and $v \in \mathbb{R}^n$, we have

$$\begin{aligned} \frac{\partial f}{\partial x^i}(\lambda v) &= \frac{\partial(f \circ M_\lambda \circ M_{1/\lambda})}{\partial x^i}(\lambda v) \\ &= \sum_{l=1}^n \frac{\partial(f \circ M_\lambda)}{\partial x^l}(v) \frac{\partial(x \mapsto x/\lambda)^l}{\partial x^i}(\lambda v) \\ &= \frac{\partial(f \circ M_\lambda)}{\partial x^i}(v) \frac{1}{\lambda} \\ &= \lambda^{r-1} \frac{\partial f}{\partial x^i}(v) \end{aligned}$$

as claimed. □