



planetmath.org

Math for the people, by the people.

IBN

Canonical name	IBN
Date of creation	2013-03-22 14:51:45
Last modified on	2013-03-22 14:51:45
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	12
Author	CWoo (3771)
Entry type	Definition
Classification	msc 16P99
Synonym	invariant basis number
Synonym	invariant dimension property
Related topic	ExampleOfFreeModuleWithBasesOfDiffrentCardinality
Defines	basis of a module
Defines	finite rank
Defines	rank of a module

## Bases of a Module

Like a vector space over a field, one can define a basis of a module  $M$  over a general ring  $R$  with 1. To simplify matter, suppose  $R$  is commutative with 1 and  $M$  is unital. A basis of  $M$  is a subset  $B = \{b_i \mid i \in I\}$  of  $M$ , where  $I$  is some ordered index set, such that every element  $m \in M$  can be uniquely written as a linear combination of elements from  $B$ :

$$m = \sum_{i \in I} r_i b_i$$

such that all but a finite number of  $r_i = 0$ .

As the above example shows, the commutativity of  $R$  is not required, and  $M$  can be assumed either as a left or right module of  $R$  (in the example above, we could take  $M$  to be the left  $R$ -module).

However, unlike a vector space, a module may not have a basis. If it does, it is called a *free module*. Vector spaces are examples of free modules over fields or division rings. If a free module  $M$  (over  $R$ ) has a finite basis with cardinality  $n$ , we often write  $R^n$  as an isomorphic copy of  $M$ .

Suppose that we are given a free module  $M$  over  $R$ , and two bases  $B_1 \neq B_2$  for  $M$ , is

$$|B_1| = |B_2|?$$

We know that this is true if  $R$  is a field or even a division ring. But in general, the equality fails. Nevertheless, it is a fact that if  $B_1$  is finite, so is  $B_2$ . So the finiteness of basis in a free module  $M$  over  $R$  is preserved when we go from one basis to another. When  $M$  has a finite basis, we say that  $M$  has *finite rank* (without saying what rank is!).

Now, even if  $M$  has finite rank, the cardinality of one basis may still be different from the cardinality of another. In other words,  $R^m$  may be isomorphic to  $R^n$  without  $m$  and  $n$  being equal.

## Invariant Basis Number

A ring  $R$  is said to have *IBN*, or *invariant basis number* if whenever  $R^m \cong R^n$  where  $m, n < \infty$ ,  $m = n$ . The positive integer  $n$  in this case is called the *rank* of module  $M$ . To rephrase, when  $F$  is a free  $R$ -module of finite rank, then  $R$  has IBN iff  $F$  has unique finite rank. Also,  $R$  has IBN iff all finite dimensional invertible matrices over  $R$  are square matrices.

### Examples

1. If  $R$  is commutative, then  $R$  has IBN.
2. If  $R$  is a division ring, then  $R$  has IBN.
3. An example of a ring  $R$  not having IBN can be found as follows: let  $V$  be a countably infinite dimensional vector space over a field. Let  $R$  be the endomorphism ring over  $V$ . Then  $R = R \oplus R$  and thus  $R^m = R^n$  for any pairs of  $(m, n)$ .