



planetmath.org

Math for the people, by the people.

## dense ring of linear transformations

Canonical name	DenseRingOfLinearTransformations
Date of creation	2013-03-22 15:38:46
Last modified on	2013-03-22 15:38:46
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	10
Author	CWoo (3771)
Entry type	Definition
Classification	msc 16K40
Synonym	dense ring
Related topic	SchursLemma
Defines	Jacobson Density Theorem

Let  $D$  be a division ring and  $V$  a vector space over  $D$ . Let  $R$  be a subring of the ring of endomorphisms (linear transformations)  $\text{End}_D(V)$  of  $V$ . Then  $R$  is said to be a *dense ring of linear transformations* (over  $D$ ) if we are given

1. any positive integer  $n$ ,
2. any set  $\{v_1, \dots, v_n\}$  of linearly independent vectors in  $V$ , and
3. any set  $\{w_1, \dots, w_n\}$  of arbitrary vectors in  $V$ ,

then there exists an element  $f \in R$  such that

$$f(v_i) = w_i \quad \text{for } i = 1, \dots, n.$$

Note that the linear independence of the  $v_i$ 's is essential in insuring the existence of a linear transformation  $f$ . Otherwise, suppose  $0 = \sum d_i v_i$  where  $d_1 \neq 0$ . Pick  $w_i$ 's so that they are linearly independent. Then  $0 = f(\sum d_i v_i) = \sum d_i f(v_i) = \sum d_i w_i$ , contradicting the linear independence of the  $w_i$ 's.

The notion of “dense” comes from topology: if  $V$  is given the discrete topology and  $\text{End}_D(V)$  the compact-open topology, then  $R$  is dense in  $\text{End}_D(V)$  iff  $R$  is a dense ring of linear transformations of  $V$ .

*Proof.* First, assume that  $R$  is a dense ring of linear transformations of  $V$ . Recall that the compact-open topology on  $\text{End}_D(V)$  has subbasis of the form  $B(K, U) := \{f \mid f(K) \subseteq U\}$ , where  $U$  is open and  $K$  is compact in  $V$ . Since  $V$  is discrete,  $K$  is finite. Now, pick a point  $g \in \text{End}_D(V)$  and let

$$B = \bigcup_{\alpha \in I} \bigcap_{i=1}^{n(\alpha)} B(K_{i\alpha}, U_{i\alpha})$$

be a neighborhood of  $g$ ,  $I$  some index set. Then for some  $\alpha \in I$ ,  $g \in \bigcap B(K_{i\alpha}, U_{i\alpha})$ . This means that  $g(K_{i\alpha}) \subseteq U_{i\alpha}$  for all  $i = 1, \dots, n(\alpha)$ . Since each  $K_{i\alpha}$  is finite, so is  $K := \bigcup K_{i\alpha}$ . After some re-indexing, let  $\{v_1, \dots, v_n\}$  be a maximal linearly independent subset of  $K$ . Set  $w_j = g(v_j)$ ,  $j = 1, \dots, n$ . By assumption, there is an  $f \in R$  such that  $f(v_j) = w_j$ , for all  $j$ . For any  $v \in K$ ,  $v$  is a linear combination of the  $v_j$ 's:  $v = \sum d_j v_j$ ,  $d_j \in D$ . Then  $f(v) = \sum d_j f(v_j) = \sum d_j w_j = g(v) \in U_{i\alpha}$  for some  $i$ . This shows that  $f(K_{i\alpha}) \subseteq U_{i\alpha}$  and we have  $f \in \bigcap B(K_{i\alpha}, U_{i\alpha}) \subseteq B$ .

Conversely, assume that the ring  $R$  is a dense subset of the space  $\text{End}_D(V)$ . Let  $v_1, \dots, v_n$  be linearly independent, and  $w_1, \dots, w_n$  be arbitrary vectors in  $V$ . Let  $W$  be the subspace spanned by the  $v_i$ 's. Because the  $v_i$ 's are linearly independent, there exists a linear transformation  $g$  such that  $g(v_i) = w_i$  and  $g(v) = 0$  for  $v \notin W$ . Let  $K_i = \{v_i\}$  and  $U_i = \{w_i\}$ . Then the  $K_i$ 's are compact and the  $U_i$ 's are open in the discrete space  $V$ . Clearly  $g \in \{h \mid h(v_i) = w_i\} = B(K_i, U_i)$  for each  $i = 1, \dots, n$ . So  $g$  lies in the neighborhood  $B = \cap B(K_i, U_i) \subseteq \text{End}_D(V)$ . Since  $R$  is dense in  $\text{End}_D(V)$ , there is an  $f \in R \cap B$ . This implies that  $f(v_i) = w_i$  for all  $i$ .  $\square$

**Remarks.**

- If  $V$  is finite dimensional over  $D$ , then any dense ring of linear transformations  $R = \text{End}_D(V)$ . This can be easily observed by using the second half of the proof above. Take a basis  $v_1, \dots, v_n$  of  $V$  and any set of  $n$  vectors  $w_1, \dots, w_n$  in  $V$ . Let  $g$  be the linear transformation that maps  $v_i$  to  $w_i$ . The above proof shows that there is an  $f \in R$  such that  $f$  agrees with  $g$  on the basis elements. But then they must agree on all of  $V$  as a result, which is precisely the statement that  $g = f \in R$ .
- It can be shown that a ring  $R$  is a primitive ring iff it is isomorphic to a dense ring of linear transformations of a vector space over a division ring. This is known as the . It is a generalization of the special case of the Wedderburn-Artin Theorem when the ring in question is a simple Artinian ring. In the general case, the finite chain condition is dropped.