



Math for the people, by the people.

the multiplicative identity of a cyclic ring
must be a generator

Canonical name	TheMultiplicativeIdentityOfACyclicRingMustBeAGenerator
Date of creation	2013-03-22 15:56:59
Last modified on	2013-03-22 15:56:59
Owner	Wkbj79 (1863)
Last modified by	Wkbj79 (1863)
Numerical id	16
Author	Wkbj79 (1863)
Entry type	Theorem
Classification	msc 16U99
Classification	msc 13F10
Classification	msc 13A99
Related topic	CyclicRing3
Related topic	CriterionForCyclicRingsToBePrincipalIdealRings
Related topic	CyclicRingsOfBehaviorOne

Theorem. *Let R be a cyclic ring with multiplicative identity u . Then u <http://planetmath.org/Generator> generates the additive group of R .*

Proof. Let k be the behavior of R . Then there exists a <http://planetmath.org/Generator> r of the additive group of R such that $r^2 = kr$. Let $a \in \mathbb{Z}$ with $u = ar$. Then $r = ur = (ar)r = ar^2 = a(kr) = (ak)r$. If R is infinite, then $ak = 1$, causing $a = k = 1$ since k is a nonnegative integer. If R is finite, then $ak \equiv 1 \pmod{|R|}$. Thus, $\gcd(k, |R|) = 1$. Since k divides $|R|$, $k = 1$. Therefore, $a \equiv 1 \pmod{|R|}$. In either case, $u = r$. \square

Note that it was also proven that, if a cyclic ring has a multiplicative identity, then it has behavior one. Its converse is also true. See <http://planetmath.org/CyclicRings0> theorem for more details.