



planetmath.org

Math for the people, by the people.

**minus one times an element is the additive  
inverse in a ring**

Canonical name	MinusOneTimesAnElementIsTheAdditiveInverseInARing
Date of creation	2013-03-22 14:14:00
Last modified on	2013-03-22 14:14:00
Owner	alozano (2414)
Last modified by	alozano (2414)
Numerical id	9
Author	alozano (2414)
Entry type	Theorem
Classification	msc 16-00
Classification	msc 13-00
Classification	msc 20-00
Synonym	$(-1) \cdot a = -a$
Related topic	0cdotA0

**Lemma 1.** *Let  $R$  be a ring (with unity 1) and let  $a$  be an element of  $R$ . Then*

$$(-1) \cdot a = -a$$

*where  $-1$  is the additive inverse of 1 and  $-a$  is the additive inverse of  $a$ .*

*Proof.* Note that for any  $a$  in  $R$  there exists a unique “ $-a$ ” by the uniqueness of additive inverse in a ring. We check that  $(-1) \cdot a$  equals the additive inverse of  $a$ .

$$\begin{aligned} a + (-1) \cdot a &= 1 \cdot a + (-1) \cdot a, && \text{by the definition of 1} \\ &= (1 + (-1)) \cdot a, && \text{by the distributive law} \\ &= 0 \cdot a, && \text{by the definition of } -1 \\ &= 0, && \text{as a result of the properties of zero} \end{aligned}$$

Hence  $(-1) \cdot a$  is “an” additive inverse for  $a$ , and by uniqueness  $(-1) \cdot a = -a$ , the additive inverse of  $a$ . Analogously, we can prove that  $a \cdot (-1) = -a$  as well.  $\square$