

involutory ring

Canonical name InvolutoryRing
Date of creation 2013-03-22 15:41:01
Last modified on 2013-03-22 15:41:01

Owner CWoo (3771) Last modified by CWoo (3771)

Numerical id 32

Author CWoo (3771) Entry type Definition Classification msc 16W10

Synonym ring admitting an involution

Synonym involutary ring
Synonym involutive ring
Synonym ring with involution
Synonym Hermitian element
Synonym symmetric element

Synonym self-adjoint
Synonym adjoint
Synonym projection
Synonym involutive ring
Related topic HollowMatrixRings

Defines involution
Defines adjoint element
Defines self-adjoint element
Defines projection element
Defines norm element

Defines skew symmetric element

trace element

Defines \*-homomorphism Defines normal element Defines unitary element

Defines

## General Definition of a Ring with Involution

Let R be a ring. An \* on R is an anti-endomorphism whose square is the identity map. In other words, for  $a, b \in R$ :

- 1.  $(a+b)^* = a^* + b^*$ ,
- $(ab)^* = b^*a^*,$
- 3.  $a^{**} = a$

A ring admitting an involution is called an *involutory ring*.  $a^*$  is called the *adjoint* of a. By (3), a is the adjoint of  $a^*$ , so that every element of R is an adjoint.

**Remark**. Note that the traditional definition of an http://planetmath.org/Involutioninvolution on a vector space is different from the one given here. Clearly, \* is bijective, so that it is an anti-automorphism. If \* is the identity on R, then R is commutative.

**Examples.** Involutory rings occur most often in rings of endomorphisms over a module. When V is a finite dimensional vector space over a field k with a given basis  $\boldsymbol{b}$ , any linear transformation over T (to itself) can be represented by a square matrix M over k via  $\boldsymbol{b}$ . The map taking M to its transpose  $M^T$  is an involution. If k is  $\mathbb{C}$ , then the map taking M to its conjugate transpose  $\overline{M}^T$  is also an involution. In general, the composition of an isomorphism and an involution is an involution, and the composition of two involutions is an isomorphism.

## \*-Homomorphisms

Let R and S be involutory rings with involutions  $*_R$  and  $*_S$ . A \*-homomorphism  $\phi: R \to S$  is a ring homomorphism which respects involutions. More precisely,

$$\phi(a^{*_R}) = \phi(a)^{*_S}$$
, for any  $a \in R$ .

By abuse of notation, if we use \* to denote both  $*_R$  and  $*_S$ , then we see that any \*-homomorphism  $\phi$  commutes with \*:  $\phi * = *\phi$ .

## Special Elements

An element  $a \in R$  such that  $a = a^*$  is called a *self-adjoint*. A ring with involution is usually associated with a ring of square matrices over a field, as such, a self-adjoint element is sometimes called a *Hermitian element*, or a *symmetric element*. For example, for any element  $a \in R$ ,

- 1.  $aa^*$  and  $a^*a$  are both self-adjoint, the first of which is called the *norm* of a. A *norm element* b is simply an element expressible in the form  $aa^*$  for some  $a \in R$ , and we write b = n(a). If  $aa^* = a^*a$ , then a is called a *normal element*. If  $a^*$  is the multiplicative inverse of a, then a is a *unitary element*. If a is unitary, then it is normal.
- 2. With respect to addition, we can also form self-adjoint elements  $a+a^*=a^*+a$ , called the *trace* of a, for any  $a \in R$ . A *trace element* b is an element expressible as  $a+a^*$  for some  $a \in R$ , and written  $b=\operatorname{tr}(a)$ .

Let S be a subset of R, write  $S^* := \{a^* \mid a \in S\}$ . Then S is said to be self-adjoint if  $S = S^*$ .

A self-adjoint that is also an idempotent in R is called a *projection*. If e and f are two projections in R such that eR = fR (principal ideals generated by e and f are equal), then e = f. For if ea = ff = f for some  $a \in R$ , then f = ea = eea = ef. Similarly, e = fe. Therefore,  $e = e^* = (fe)^* = e^*f^* = ef = f$ .

If the characteristic of R is not 2, we also have a companion concept to self-adjointness, that of skew symmetry. An element a in R is skew symmetric if  $a = -a^*$ . Again, the name of this is borrowed from linear algebra.