



planetmath.org

Math for the people, by the people.

octonion

Canonical name	Octonion
Date of creation	2013-03-22 15:21:42
Last modified on	2013-03-22 15:21:42
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	15
Author	CWoo (3771)
Entry type	Definition
Classification	msc 17A75
Classification	msc 17D05
Synonym	Cayley algebra
Related topic	TheoremsOnSumsOfSquares
Related topic	DivisionAlgebra
Defines	octonion algebra

Let \mathbb{H} be the quaternions over the reals \mathbb{R} . Apply the Cayley-Dickson construction to \mathbb{H} once, and we obtain an algebra, variously called *Cayley algebra*, *the octonion algebra*, or simply *the octonions*, over \mathbb{R} . Specifically the construction is carried out as follows:

1. Form the vector space $\mathbb{O} = \mathbb{H} \oplus \mathbb{H}\mathbf{k}$; any element of \mathbb{O} can be written as $a + b\mathbf{k}$, where $a, b \in \mathbb{H}$;
2. Define a binary operation on \mathbb{O} called *the multiplication in \mathbb{O}* by

$$(a + b\mathbf{k})(c + d\mathbf{k}) := (ac - \bar{d}b) + (da + b\bar{c})\mathbf{k},$$

where $a, b, c, d \in \mathbb{H}$, and \bar{c} is the quaternionic conjugation of $c \in \mathbb{H}$. When $b = d = 0$, the multiplication is reduced the multiplication in \mathbb{H} . In addition, the multiplication rule above imply the following:

$$a(d\mathbf{k}) = (da)\mathbf{k} \tag{1}$$

$$(b\mathbf{k})c = (b\bar{c})\mathbf{k} \tag{2}$$

$$(b\mathbf{k})(d\mathbf{k}) = -\bar{d}b. \tag{3}$$

In particular, in the last equation, if $b = d = 1$, $\mathbf{k}^2 = -1$.

3. Define a unary operation on \mathbb{O} called *the octonionic conjugation in \mathbb{O}* by

$$\overline{a + b\mathbf{k}} := \bar{a} - b\mathbf{k},$$

where $a, b \in \mathbb{H}$. Clearly, the octonionic conjugation is an <http://planetmath.org/Involution> ($\overline{\bar{x}} = x$).

4. Finally, define a unary operation N on \mathbb{O} called *the norm in \mathbb{O}* by $N(x) := x\bar{x}$, where $x \in \mathbb{O}$. Write $x = a + b\mathbf{k}$, then

$$N(x) = (a + b\mathbf{k})(\bar{a} - b\mathbf{k}) = (a\bar{a} + \bar{b}b) + (-ba + b\bar{\bar{a}})\mathbf{k} = a\bar{a} + b\bar{b} \geq 0.$$

It is not hard to see that $N(x) = 0$ iff $x = 0$.

The above four (actually, only the first two suffice) steps makes \mathbb{O} into an 8-dimensional algebra over \mathbb{R} such that \mathbb{H} is embedded as a subalgebra.

With the last two steps, one can define the inverse of a non-zero element $x \in \mathbb{O}$ by

$$x^{-1} := \frac{\bar{x}}{N(x)}$$

so that $xx^{-1} = x^{-1}x = 1$. Since x is arbitrary, \mathbb{O} has no zero divisors. Upon checking that $x^{-1}(xy) = y = (yx)x^{-1}$, the non-associative algebra \mathbb{O} is turned into a division algebra.

Since $N(x) \geq 0$ for any $x \in \mathbb{O}$, we can define a non-negative real-valued function $\|\cdot\|$ on \mathbb{O} by $\|x\| = \sqrt{N(x)}$. This is clearly well-defined and $\|x\| = 0$ iff $x = 0$. In addition, it is not hard to see that, for any $r \in \mathbb{R}$ and $x \in \mathbb{O}$, $\|rx\| = |r|\|x\|$, and that $\|\cdot\|$ satisfies the triangular inequality. This makes \mathbb{O} into a normed division algebra.

Since the multiplication in \mathbb{H} is noncommutative, \mathbb{O} is noncommutative. In fact, if we write $\mathbb{H} = \mathbb{C} \oplus \mathbb{C}\mathbf{j}$, where \mathbb{C} are the complex numbers and $\mathbf{j}^2 = -1$, then $B = \{1, \mathbf{i}, \mathbf{j}, \mathbf{ij}\}$ is a basis for the vector space \mathbb{H} over \mathbb{R} . With the introduction of $\mathbf{k} \in \mathbb{O}$, we quickly check that \mathbf{k} anti-commute with the non-real basis elements in B :

$$\mathbf{ik} = -\mathbf{ki}, \quad \mathbf{jk} = -\mathbf{kj}, \quad (\mathbf{ij})\mathbf{k} = -\mathbf{k}(\mathbf{ij}).$$

Furthermore, one checks that $\mathbf{i}(\mathbf{jk}) = (\mathbf{ji})\mathbf{k} = -(\mathbf{ij})\mathbf{k}$, so that \mathbb{O} is not associative.

Since $\mathbb{O} = \mathbb{H} \oplus \mathbb{H}\mathbf{k}$, the set $\{1, \mathbf{i}, \mathbf{j}, \mathbf{ij}, \mathbf{k}, \mathbf{ik}, \mathbf{jk}, (\mathbf{ij})\mathbf{k}\} (= B \cup B\mathbf{k})$ is a basis for \mathbb{O} over \mathbb{R} . A less messy way to represent these basis elements is done the following assignment:

basis element	1	\mathbf{i}	\mathbf{j}	\mathbf{ij}	\mathbf{k}	\mathbf{ik}	\mathbf{jk}	$(\mathbf{ij})\mathbf{k}$
basis element rewritten	\mathbf{i}_0	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6	\mathbf{i}_7

Any element x of \mathbb{O} can thus be expressed uniquely as $\sum_{n=0}^7 r_n \mathbf{i}_n$, where $r_n \in \mathbb{R}$. Using Equations (1)-(3) above, one can form a multiplication table for these basis elements \mathbf{i}_n 's:

row \times column	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3	\mathbf{i}_4	\mathbf{i}_5	\mathbf{i}_6	\mathbf{i}_7
\mathbf{i}_1	-1	\mathbf{i}_3	$-\mathbf{i}_2$	\mathbf{i}_5	$-\mathbf{i}_4$	$-\mathbf{i}_7$	\mathbf{i}_6
\mathbf{i}_2	$-\mathbf{i}_3$	-1	\mathbf{i}_1	\mathbf{i}_6	\mathbf{i}_7	$-\mathbf{i}_4$	$-\mathbf{i}_5$
\mathbf{i}_3	\mathbf{i}_2	$-\mathbf{i}_1$	-1	\mathbf{i}_7	$-\mathbf{i}_6$	\mathbf{i}_5	$-\mathbf{i}_4$
\mathbf{i}_4	$-\mathbf{i}_5$	$-\mathbf{i}_6$	$-\mathbf{i}_7$	-1	\mathbf{i}_1	\mathbf{i}_2	\mathbf{i}_3
\mathbf{i}_5	\mathbf{i}_4	$-\mathbf{i}_7$	\mathbf{i}_6	$-\mathbf{i}_1$	-1	$-\mathbf{i}_3$	\mathbf{i}_2
\mathbf{i}_6	\mathbf{i}_7	\mathbf{i}_4	$-\mathbf{i}_5$	$-\mathbf{i}_2$	\mathbf{i}_3	-1	$-\mathbf{i}_1$
\mathbf{i}_7	$-\mathbf{i}_6$	\mathbf{i}_5	\mathbf{i}_4	$-\mathbf{i}_3$	$-\mathbf{i}_2$	\mathbf{i}_1	-1

Other well known properties of the octonions are

1. $\overline{xy} = \overline{y} \overline{x}$ for any $x, y \in \mathbb{O}$.
2. $N(xy) = N(x)N(y)$ so that \mathbb{O} is a composition algebra. It also proves that the product of sums of eight squares is a sum of eight squares.
3. \mathbb{O} is an alternative algebra. As a result, any two elements of \mathbb{O} generate an associative algebra. In fact, the algebra is isomorphic to one of \mathbb{R} , \mathbb{C} , and \mathbb{H} . This is the consequence of <http://planetmath.org/ArtinsTheoremOnAlternativeAlgebras> Theorem.