



planetmath.org

Math for the people, by the people.

## equivalent representations of groupoids

Canonical name	EquivalentRepresentationsOfGroupoids
Date of creation	2013-03-22 18:16:30
Last modified on	2013-03-22 18:16:30
Owner	bci1 (20947)
Last modified by	bci1 (20947)
Numerical id	25
Author	bci1 (20947)
Entry type	Definition
Classification	msc 18D05
Classification	msc 55N33
Classification	msc 55N20
Classification	msc 55P10
Classification	msc 55U40
Classification	msc 55U99
Synonym	equivalence class of groupoid representations
Related topic	Groupoids
Related topic	CAlgebra3
Related topic	QuantumOperatorAlgebrasInQuantumFieldTheories
Defines	equivalent representations of groupoids

**Definition 0.1.** Two representations of groupoids  $(\mu_i, U_{\mathbf{G}} * \mathcal{H}, L_i)$  , for  $i = 1, 2$  are called *equivalent* if  $\mu_1 \sim \mu_2$ , and if there also exists a fiber-preserving isomorphism of analytical Hilbert space bundles  $v : (U_{\mathbf{G}} * \mathcal{H}_1)|_U \longrightarrow (U_{\mathbf{G}} * \mathcal{H}_2)|_U$  , where  $U$  is a measurable subset of  $U_{\mathbf{G}}$  of null complementarity; the isomorphism  $v$  also has the following property:  $\hat{v}[r(x)]\hat{L}_1(x) = \hat{L}_2\hat{v}[d(x)]$  for  $x \in \mathbf{G}|_U$ .