



Math for the people, by the people.

Leray's theorem

Canonical name	LeraysTheorem
Date of creation	2013-03-22 14:42:32
Last modified on	2013-03-22 14:42:32
Owner	mathcam (2727)
Last modified by	mathcam (2727)
Numerical id	9
Author	mathcam (2727)
Entry type	Theorem
Classification	msc 18G60
Related topic	sheaf
Related topic	sheafcohomology
Related topic	SheafCohomology

Let \mathcal{F} be a sheaf on a topological space X and $\mathcal{U} = \{U_i\}$ an open cover of X . If \mathcal{F} is <http://planetmath.org/AcyclicSheaf> acyclic on every element of \mathcal{U} , then

$$\check{H}^q(\mathcal{U}, \mathcal{F}) = \check{H}^q(X, \mathcal{F}),$$

where $\check{H}^q(\mathcal{U}, \mathcal{F})$ is the q -th <http://planetmath.org/CechCohomologyGroup> Cech cohomology group of \mathcal{F} with respect to the open cover \mathcal{U} .

References

- [1] Bonavero, Laurent. *Cohomology of Line Bundles on Toric Varieties, Vanishing Theorems*. Lectures 16-17 from “Summer School 2000: Geometry of Toric Varieties.”