



planetmath.org

Math for the people, by the people.

quantum topos

Canonical name	QuantumTopos
Date of creation	2013-03-22 18:17:48
Last modified on	2013-03-22 18:17:48
Owner	bci1 (20947)
Last modified by	bci1 (20947)
Numerical id	25
Author	bci1 (20947)
Entry type	Definition
Classification	msc 18D25
Classification	msc 18-00
Classification	msc 55U99
Classification	msc 81-00
Classification	msc 81P05
Classification	msc 81Q05
Synonym	quantum category
Related topic	NonCommutativeStructureAndOperation
Related topic	Commutative
Related topic	QuantumCategory
Related topic	HeytingAlgebra
Related topic	AxiomsOfTopoi
Related topic	QuantumLogic
Related topic	CategoricalDynamics
Related topic	Lattice
Related topic	IntuitionisticLogic
Related topic	ConsistencyOfClassicNumberTheory
Defines	quantum state space

Preliminary Data.

There are several distinct definitions of *quantum topos* in the Mathematical Physics literature attempting to redefine the quantum logic that was first introduced by von Neumann and Birkhoff for the foundation of Quantum Mechanics. The definitions of quantum topoi published so far are not, however, those of ‘quantum’ categories (previously introduced as rigid monoidal categories) - with finite limits and power objects.

Definition 0.1. A *quantum topos* was defined as a *general model, or representation of quantum state spaces (QST) in a topos* with a <http://planetmath.org/Commutative>(commutative) Heyting logic algebra as a subobject (*quantum logic*) classifier. The differences between the several published definitions of a quantum topos differ in the categorical representation in QST’s, and in the choice of category, but not in the choice of quantum logic algebra that was selected as a standard, Heyting logic algebra (or *Heyting algebra*) which has a commutative *Heyting lattice* structure; this choice is at variance with the original quantum logic introduced by von Neumann and Birkhoff. Thus instead of the orthomodular lattice of Birkhoff and von Neumann, the recent definitions of quantum topoi postulate an intuitionistic- Brouwer logic corresponding to a pseudocomplemented and rel. pseudocomplemented lattice structure, as further explained in the next section.

0.1 Heyting Logic Concept and Algebraic Structure

Definition 0.2. A *Heyting lattice* L is a Brouwer-intuitionistic logic lattice with a bottom, or lowest element 0. In the more technical classification it is a <http://planetmath.org/Commutative> commutative lattice which is both ‘pseudocomplemented and also relatively pseudocomplemented’. The concept of <http://planetmath.org/RelativelyPseudocomplemented> relative pseudocomplementation coincides with the material implication operator, \Rightarrow , in symbolic propositional logic based on chrysippian or Boolean logic.

Definition 0.3. A *Heyting algebra* is a p -algebra (as defined next in **Definition 1.3**) with the relative pseudocomplementation operation \rightarrow (which replaces the propositional implication \Rightarrow).

Given an element a in a bounded lattice L , a *complement* of a is defined to be an element $b \in L$, if such an element exists, such that

$$a \wedge b = 0, \quad \text{and} \quad a \vee b = 1.$$

To surmount the non-uniqueness of the complement, an alternative to the latter was defined—the *pseudocomplement* of an element.

An element b in a lattice L with 0 is a *pseudocomplement* of $a \in L$ if

1. $b \wedge a = 0$
2. for any c such that $c \wedge a = 0$ then $c \leq b$.

In other words, b is the maximal element in the set $\{c \in L \mid c \wedge a = 0\}$.

Definition 0.4. A convenient modification of the pseudocomplemented (pc) lattice concept is a *p-algebra* (or pseudocomplemented algebra) which is a pc-lattice where $*$ is regarded as an algebraic operator. Thus, a morphism of pc-lattices is a proper lattice homomorphism, whereas a morphism between two p-algebras is a lattice homomorphism f that also preserves the pc-algebraic operation $*$, i.e., $f(a^*) = f(a)^*$. One can therefore define a *category of p-algebras* by specifying the morphism between any pair of p-algebras (considered as objects of this algebraic logic category) as the $\{0, 1\}$ -lattice homomorphism, with the following condition $f(1) = f(0^*) = f(0)^* = 0^* = 1$ being also satisfied.

Remark Unlike the Heyting lattice, an LM_n -logic algebra has a *non-commutative* lattice structure and is therefore considered as a stronger candidate for quantum logics, including those based on the orthomodular lattices of the original quantum logic of Birkhoff and von Neumann. Thus, a generalized topos defined with a subobject classifier based on LM_n -logic algebra may provide suitable representations of arbitrary quantum state spaces.

References

Butterfield, J. and C. J. Isham: 2001, Space-time and the Philosophical Challenges of Quantum Gravity., in C. Callender and N. Hugget (eds.) *Physics Meets Philosophy at the Planck scale.*, Cambridge University Press, pp.33–89. Butterfield, J. and C. J. Isham: 1998, 1999, 2000–2002, A topos perspective on the Kochen–Specker theorem I - IV, *Int. J. Theor. Phys.* **37** No 11., 2669–2733 **38** No 3., 827–859, **39** No 6., 1413–1436, **41** No 4., 613–639.