



planetmath.org

Math for the people, by the people.

topological groupoid

Canonical name	TopologicalGroupoid
Date of creation	2013-03-22 14:35:11
Last modified on	2013-03-22 14:35:11
Owner	HkBst (6197)
Last modified by	HkBst (6197)
Numerical id	20
Author	HkBst (6197)
Entry type	Definition
Classification	msc 18B40
Classification	msc 20L05
Classification	msc 54H13
Classification	msc 54H11
Related topic	Groupoids
Related topic	LocallyCompactGroupoids
Defines	groupoid
Defines	transitive groupoid
Defines	principal groupoid
Defines	isotropy group
Defines	topological groupoid
Defines	domain map
Defines	range map
Defines	unit space
Defines	isotropy group

A *groupoid* is a set  $G$  together with a subset  $G_2 \subset G^2$  of composable pairs, a *multiplication*  $\mu : G_2 \rightarrow G : (a, b) \mapsto ab$  and an *inversion*  $\cdot^{-1} : G \rightarrow G : a \mapsto a^{-1}$  such that

1.  $\cdot^{-1} \circ \cdot^{-1} = \text{id}_G$ ,
2. if  $\{(a, b), (b, c)\} \subset G_2$  then  $\{(ab, c), (a, bc)\} \subset G_2$  and  $(ab)c = a(bc)$ ,
3.  $(b, b^{-1}) \in G_2$  and if  $(a, b) \in G_2$  then  $abb^{-1} = a$  and
4.  $(b^{-1}, b) \in G_2$  and if  $(b, c) \in G_2$  then  $b^{-1}bc = c$ .

Furthermore we have the *source* or *domain map*  $\sigma : G \rightarrow G : a \mapsto a^{-1}a$  and the *target* or *range map*  $\tau : G \rightarrow G : a \mapsto aa^{-1}$ . The image of these maps is called the *unit space* and denoted  $G_0$ . If the unit space is a singleton then we regain the notion of a group.

We also define  $G_a := \sigma^{-1}(\{a\})$ ,  $G^b := \tau^{-1}(\{b\})$  and  $G_a^b := G_a \cap G^b$ . It is not hard to see that  $G_a^a$  is a group, which is called the *isotropy group* at  $a$ .

We say that a groupoid  $G$  is *principal* and *transitive*, if the map  $(\sigma, \tau) : G \rightarrow G_0 \times G_0$  is injective and surjective, respectively.

A groupoid can be more abstractly and more succinctly defined as a category whose morphisms are all isomorphisms.

A *topological groupoid* is a groupoid  $G$  which is also a topological space, such that the multiplication and inversion are continuous when  $G_2$  is endowed with the induced product topology from  $G^2$ . Consequently also  $\sigma$  and  $\tau$  are continuous.

## References

- [1] P.J. Higgins, *Categories and groupoids*, van Nostrand original, 1971; Reprint Theory and Applications of Categories, 7 (2005) pp 1-195.
- [2] R. Brown, *Topology and groupoids*, xxv+512pp, Booksurge 2006.
- [3] R. Brown, ‘Three themes in the work of Charles Ehresmann: Local-to-global; Groupoids; Higher dimensions’, *Proceedings of the 7th Conference on the Geometry and Topology of Manifolds: The Mathematical Legacy of Charles Ehresmann, Bedlewo (Poland) 8.05.2005-15.05.2005*, Banach Centre Publications 76, Institute of Mathematics Polish Academy of Sciences, Warsaw, (2007) 51-63. (math.DG/0602499).