



planetmath.org

Math for the people, by the people.

groupoid (category theoretic)

Canonical name	GroupoidcategoryTheoretic
Date of creation	2013-03-22 13:08:17
Last modified on	2013-03-22 13:08:17
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	19
Author	CWoo (3771)
Entry type	Definition
Classification	msc 18B40
Classification	msc 20L05
Synonym	groupoid
Synonym	virtual group
Related topic	BrandtGroupoid
Defines	composable pair

A *groupoid*, also known as a *virtual group*, is a small category where every morphism is invertible. We can give a more explicit, algebraic definition: start with a set G , and a partial binary operation \circ on G . Call a pair (x, y) of elements of G a *composable pair* if $(x, y) \in \text{dom}(\circ)$. A *groupoid* is the pair (G, \circ) , together with two unary operations e_L and e_R on it, satisfying the following conditions:

1. (x, y) is a composable pair iff $e_R(x) = e_L(y)$.
2. (x, y) and $(x \circ y, z)$ are composable pairs iff (y, z) and $(x, y \circ z)$ are, and if one of these is true, then $(x \circ y) \circ z = x \circ (y \circ z)$.
3. $(e_L(x), x)$ and $(x, e_R(x))$ are composable pairs and $x = e_L(x) \circ x = x \circ e_R(x)$.
4. for each $x \in G$, there exists $y \in G$ such that (x, y) and (y, x) are composable pairs, and $e_L(x) = x \circ y$ and $e_R(x) = y \circ x$.

Below are some properties:

1. In condition 4 above, $e_L(x) = e_R(y)$ and $e_R(x) = e_L(y)$. This is true by condition 1, since both (x, y) and (y, x) are composable pairs.
2. Again, in condition 4, y is unique. To see this, suppose $z \in G$ satisfies condition 4 (in place of y). Then $y = y \circ e_R(y) = y \circ e_L(x) = y \circ (x \circ y) = y \circ (x \circ z) = (y \circ x) \circ z = (z \circ x) \circ z = e_R(x) \circ z = e_L(z) \circ z = z$. Notice property 1 is used in the proof. We call y the *inverse* of x , and write x^{-1} .
3. In view of condition 4, both e_L and e_R are unique. In other words, if $f_L, f_R : G \rightarrow G$ are unary operators on G satisfying conditions 3 and 4 above (in place of e_L and e_R), then $f_L = e_L$ and $f_R = e_R$. In fact, $e_L(x) = x \circ x^{-1}$ and $e_R(x) = x^{-1} \circ x$.
4. Since $x = e_L(x) \circ x = e_L(x) \circ (e_L(x) \circ x) = (e_L(x) \circ e_L(x)) \circ x$, we see that $e_L(x)$ is composable with itself, and that $e_L(x) \circ e_L(x) = e_L(x)$ by the previous property. Similarly, $e_R(x) \circ e_R(x) = e_R(x)$. This shows that $e_R(x)$ and $e_L(x)$ are idempotent with respect to \circ for every $x \in G$.
5. Since $(e_L(x), x)$ is a composable pair, $e_R(e_L(x)) = e_L(x)$ for any $x \in G$. Similarly, $e_L(e_R(x)) = e_R(x)$. Hence $e_R(e_R(x)) = e_R(e_L(e_R(x))) =$

$e_L(e_R(x)) = e_R(x)$. Similarly, $e_L(e_L(x)) = e_L(x)$. This shows that e_R and e_L are idempotent with respect to functional compositions.

6. (Cancellation property): if $x \circ y = x \circ z$, then $y = z$; if $y \circ x = z \circ x$, then $y = z$.

Proof. Since (x, y) is a composable pair, $e_R(x) = e_L(y)$. But $e_R(e_R(x)) = e_R(x)$, we have $e_R(e_R(x)) = e_L(y)$ so that $(e_R(x), y) = (x^{-1} \circ x, y)$ is a composable pair, hence $(x^{-1}, x \circ y)$ is a composable pair and $x^{-1} \circ (x \circ y) = (x^{-1} \circ x) \circ y = e_R(x) \circ y$. Since $(e_R(x), y)$ is a composable pair, $e_R(x) = e_R(e_R(x)) = e_L(y)$. As a result, $x^{-1} \circ (x \circ y) = e_L(y) \circ y = y$. Similarly $x^{-1} \circ (x \circ z) = z$. By assumption, we deduce that $y = z$. The other statement is proved similarly. \square

7. The algebraic definition given can be easily turned into a categorical definition (using objects and morphisms). The details are left for the reader.

If e_R and e_L are constant functions, then G is a group.

Remark. There is also a <http://planetmath.org/Groupoidgroup-theoretic> concept with the same name.