



planetmath.org

Math for the people, by the people.

direct limit of sets

Canonical name	DirectLimitOfSets
Date of creation	2013-03-22 16:52:36
Last modified on	2013-03-22 16:52:36
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	11
Author	CWoo (3771)
Entry type	Example
Classification	msc 18A30
Synonym	direct system
Synonym	inverse system
Synonym	projective system
Related topic	DirectLimitOfAlgebraicSystems
Defines	direct family
Defines	inverse family
Defines	inverse limit of sets

Let  $\mathcal{A} = \{A_i \mid i \in I\}$  be a family of sets indexed by a non-empty set  $I$ .  $\mathcal{A}$  is said to be a *direct family* if

1.  $I$  is a directed set,
2. whenever  $i \leq j$  in  $I$ , there is a function  $\phi_{ij} : A_i \rightarrow A_j$ ,
3.  $\phi_{ii}$  is the identity function on  $A_i$ ,
4. if  $i \leq j \leq k$ , then  $\phi_{jk} \circ \phi_{ij} = \phi_{ik}$ .

In the last condition, if we write  $a\phi_{ij} := \phi_{ij}(a)$  for  $a \in A_i$ , then the equation can be rewritten as  $\phi_{ij}\phi_{jk} = \phi_{ik}$ .

For example, the natural numbers  $\mathbb{N} = \{1, 2, \dots, n, \dots\}$  can be regarded as a direct family. Here, for any  $i \leq j$ ,  $\phi_{ij} : i \rightarrow j$  is given by the natural injection  $\phi_{ij}(\ell) := \ell$  for any  $\ell \in i$ .

Let  $\mathcal{A}$  be a direct family of sets, indexed by  $I$ . Take the disjoint union of the members of  $\mathcal{A}$  and call it  $A$  (this can be achieved even when the members themselves have non-empty intersections, simply form the product  $A_i \times \{i\}$  first before taking the union). Therefore,  $A$  has the properties that

- for any  $a \in A$ ,  $a \in A_i$  for some  $i \in I$ , and
- if  $a \in A_i$  and  $b \in A_j$  and  $i \neq j$ , then  $a \neq b$ .

Define a binary relation  $\sim$  on  $A$  as follows: given that  $a \in A_i$  and  $b \in A_j$ ,  $a \sim b$  iff there is  $A_k$  such that  $\phi_{ik}(a) = \phi_{jk}(b)$ .

**Proposition 1.**  $\sim$  on  $A$  is an equivalence relation.

*Proof.* Clearly,  $\sim$  is symmetric. By condition 2 of a direct family,  $\sim$  is also reflexive. Now, suppose  $a \sim b$  and  $b \sim c$  with  $a \in A_i$ ,  $b \in A_j$  and  $c \in A_k$ . So there are  $p, q \in I$  such that  $\phi_{ip}(a) = \phi_{jp}(b)$  and  $\phi_{jq}(b) = \phi_{kq}(c)$ . Since  $I$  is directed, there is  $r \in I$  such that  $p, q \leq r$ . From this, we have  $\phi_{ir}(a) = \phi_{pr}(\phi_{ip}(a)) = \phi_{pr}(\phi_{jp}(b)) = \phi_{jr}(b)$ . Similarly,  $\phi_{kr}(c) = \phi_{qr}(\phi_{kq}(c)) = \phi_{qr}(\phi_{jq}(b))$ . Hence  $a \sim c$ .  $\square$

**Definition.** Let  $\mathcal{A}$  be a direct family of sets indexed by  $I$ . Let  $A$  and  $\sim$  be defined as above. Then the quotient  $A/\sim$  is called the *direct limit* of the sets in  $\mathcal{A}$ . The direct limit of sets  $A_i$  is sometimes written  $A_\infty$ , or  $\varinjlim A_i$ . Elements of  $A_\infty$  are sometimes denoted by  $[a]_I$  or  $[a]$  whenever there is no confusion.

**Remarks.**

- This definition is consistent with the formal definition of direct limits in a category. The index  $I$ , being a directed set, can be viewed as a category whose objects are elements of  $I$  and morphisms defined by the partial order on  $I$ .
- The notation  $A_\infty$  comes from the following fact: if  $|I| = n < \infty$ , then  $\varinjlim A_i \cong A_n$ . Here,  $\cong$  stands for bijection.
- For every  $i \in I$ , there is a natural mapping  $A_i \rightarrow A_\infty$ , given by  $a \mapsto [a]_I$ . This map may be variously denoted by  $\phi_i$ ,  $\phi_{i\infty}$ , or  $\phi_{iI}$ .
- Let  $J$  be a subset of a directed set  $I$ . Let  $\mathcal{A}$  be a direct family indexed by  $I$  and  $\mathcal{A}' \subseteq \mathcal{A}$  indexed by  $J$ . Form the direct limit  $A'_\infty$  of sets in  $\mathcal{A}'$ . Then there is a natural mapping  $\phi_{JI} : A'_\infty \rightarrow A_\infty$  such that for any  $j \in J$ ,  $\phi_{JI} \circ \phi_{jJ} = \phi_{jI}$ .

The dual notion of a direct limit of sets is that of an inverse limit. Instead of starting from a direct family of sets, we start with an *inverse family* of sets, which is defined similarly to that of a direct family, except  $I$  is a filtered set, and the mappings  $\phi_{ij} : A_i \rightarrow A_j$  is defined whenever  $j \leq i$ . An inverse family is also known as an *inverse system*, or a *projective system*.