



planetmath.org

Math for the people, by the people.

superalgebroids and higher dimensional  
algebroids

Canonical name	SuperalgebroidsAndHigherDimensionalAlgebroids
Date of creation	2013-03-22 18:30:02
Last modified on	2013-03-22 18:30:02
Owner	bci1 (20947)
Last modified by	bci1 (20947)
Numerical id	10
Author	bci1 (20947)
Entry type	Topic
Classification	msc 18A25
Classification	msc 18A20
Classification	msc 18A15
Classification	msc 55U40
Classification	msc 55U35
Classification	msc 18D05
Synonym	HDA superstructure
Related topic	RAlgebroid
Related topic	HDA
Related topic	HigherDimensionalAlgebraHDA
Related topic	SuperfieldsSuperspace
Defines	generalized superalgebra
Defines	graded Lie algebroid
Defines	HDA structure

### 0.0.1 Definitions of double, and higher dimensional, algebroids, superalgebroids and generalized superalgebras

#### Double algebroids

**Definition 0.1.** A *double  $R$ -algebroid* consists of a <http://planetmath.org/HomotopyDoubleGroupoid> category  $D$ , as detailed in ref.[?], such that each category structure has the additional structure of an  $R$ -algebroid. More precisely, a double <http://planetmath.org/RAlgebroid>  $D$  involves four related <http://planetmath.org/RAlgebroid>  $R$ -algebroids:

$$\begin{aligned} (D, D_1, \partial_1^0, \partial_1^1, \varepsilon_1, +_1, \circ_1, \cdot_1), & \quad (D, D_2, \partial_2^0, \partial_2^1, \varepsilon_2, +_2, \circ_2, \cdot_2) \\ (D_1, D_0, \delta_1^0, \delta_1^1, \varepsilon, +, \circ, \cdot), & \quad (D_2, D_0, \delta_2^0, \delta_2^1, \varepsilon, +, \circ, \cdot) \end{aligned} \quad (0.1)$$

that satisfy the following rules:

i)  $\delta_2^i \partial_2^j = \delta_1^j \partial_1^i$  for  $i, j \in \{0, 1\}$

ii)

$$\begin{aligned} \partial_2^i(\alpha +_1 \beta) &= \partial_2^i \alpha + \partial_2^i \beta, & \partial_1^i(\alpha +_2 \beta) &= \partial_1^i \alpha + \partial_1^i \beta \\ \partial_2^i(\alpha \circ_1 \beta) &= \partial_2^i \alpha \circ \partial_2^i \beta, & \partial_1^i(\alpha \circ_2 \beta) &= \partial_1^i \alpha \circ \partial_1^i \beta \end{aligned} \quad (0.2)$$

for  $i = 0, 1, \alpha, \beta \in D$  and both sides are defined.

iii)

$$\begin{aligned} r_{\cdot 1}(\alpha +_2 \beta) &= (r_{\cdot 1} \alpha) +_2 (r_{\cdot 1} \beta), & r_{\cdot 2}(\alpha +_1 \beta) &= (r_{\cdot 2} \alpha) +_1 (r_{\cdot 2} \beta) \\ r_{\cdot 1}(\alpha \circ_2 \beta) &= (r_{\cdot 1} \alpha) \circ_2 (r_{\cdot 1} \beta), & r_{\cdot 2}(\alpha \circ_1 \beta) &= (r_{\cdot 2} \alpha) \circ_1 (r_{\cdot 2} \beta) \\ r_{\cdot 1}(s_{\cdot 2} \beta) &= s_{\cdot 2}(r_{\cdot 1} \beta) \end{aligned} \quad (0.3)$$

for all  $\alpha, \beta \in D, r, s \in R$  and both sides are defined.

iv)

$$\begin{aligned} (\alpha +_1 \beta) +_2 (\gamma +_1 \lambda) &= (\alpha +_2 \gamma) +_1 (\beta +_2 \lambda), \\ (\alpha \circ_1 \beta) \circ_2 (\gamma \circ_1 \lambda) &= (\alpha \circ_2 \gamma) \circ_1 (\beta \circ_2 \lambda) \\ (\alpha +_i \beta) \circ_j (\gamma +_i \lambda) &= (\alpha \circ_j \gamma) +_i (\beta \circ_j \lambda) \end{aligned} \quad (0.4)$$

for  $i \neq j$ , whenever both sides are defined.

The definition of a double algebroid specified above was introduced by Brown and Mosa [?]. Two functors can be then constructed, one from the category of double algebroids to the category of crossed modules of algebroids, whereas the reverse functor is the unique adjoint (up to natural equivalence). The construction of such functors requires the following definition.

## 0.1 Category of Double Algebroids

A *morphism*  $f : D \rightarrow \mathcal{E}$  of double algebroids is then defined as a morphism of truncated cubical sets which commutes with all the algebroid structures. Thus, one can construct a category **DA** of double algebroids and their morphisms. The main construction in this subsection is that of two functors  $\eta, \eta'$  from this category **DA** to the category **CM** of crossed modules of algebroids.

Let  $D$  be a double algebroid. One can associate to  $D$  a crossed module  $\mu : M \rightarrow D_1$ . Here  $M(x, y)$  will consist of elements  $m$  of  $D$  with boundary of the form:  $0 \ 1$

$$\partial m = \begin{pmatrix} & a & \\ 1_y & 0_{xy} & 1_x \end{pmatrix}, \quad (0.5)$$

that is  $M(x, y) = \{m \in D : \partial_1^1 m = 0_{xy}, \partial_2^0 m = 1_x, \partial_2^1 m = 1_y\}$ .

## 0.2 Cubic and Higher dimensional algebroids

One can extend the above notion of double algebroid to cubic and higher dimensional algebroids. The concepts of 2-algebroid, 3-algebroid, ...,  $n$ -algebroid and superalgebroid are however quite distinct from those of double, cubic, ...,  $n$ -tuple algebroid, and have technically less complicated definitions.

## References

- [1] R. Brown and G. H. Mosa: Double algebroids and crossed modules of algebroids, University of Wales–Bangor, Maths Preprint, 1986.
- [2] R. Brown and C.B. Spencer: Double groupoids and crossed modules, *Cahiers Top. Géom.Diff.* **17**: 343–362 (1976).