



planetmath.org

Math for the people, by the people.

splicing together exact sequences

Canonical name	SplicingTogetherExactSequences
Date of creation	2013-03-22 19:03:35
Last modified on	2013-03-22 19:03:35
Owner	rm50 (10146)
Last modified by	rm50 (10146)
Numerical id	4
Author	rm50 (10146)
Entry type	Theorem
Classification	msc 18G35
Synonym	splicing lemma

This article proves a simple but very useful result about “splicing” together two exact sequences. Assume we are working in an abelian category such as groups, rings, or modules.

Proposition 1. *Let*

$$A \rightarrow B \xrightarrow{f} C$$

and

$$D \xrightarrow{g} E \rightarrow F$$

be exact, and assume that there is an isomorphism $\varphi : \operatorname{coker} f \rightarrow \ker g$. Define $\psi : C \rightarrow D : c \mapsto \varphi(\bar{c})$, where \bar{c} is the image of c in $\operatorname{coker} f$. Then the following is exact:

$$A \rightarrow B \xrightarrow{f} C \xrightarrow{\psi} D \xrightarrow{g} E \rightarrow F$$

Proof. Exactness at C :

$$c \in \ker \psi \iff \psi(c) = \varphi(\bar{c}) = 0 \iff \bar{c} = 0 \iff c \in \operatorname{im} f.$$

Exactness at D :

$$d \in \ker g \iff d = \varphi(\bar{c}) \text{ for some } c \in C \iff d = \psi(c) \text{ for some } c \in C.$$

□