



planetmath.org

Math for the people, by the people.

essentially surjective

Canonical name	EssentiallySurjective
Date of creation	2013-03-22 15:18:16
Last modified on	2013-03-22 15:18:16
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	13
Author	CWoo (3771)
Entry type	Definition
Classification	msc 18A22
Synonym	dense functor
Synonym	isomorphism-dense in
Synonym	isomorphism-dense
Related topic	EquivalenceOfCategories
Defines	isomorphism-dense subcategory

Let \mathcal{C} and \mathcal{D} be categories. A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is *essentially surjective* if for any object $A \in \mathcal{OB}(\mathcal{D})$, there exists an object $X \in \mathcal{OB}(\mathcal{C})$, such that $F(X) \cong A$. That is, there are morphisms (in \mathcal{D}) $f: F(X) \rightarrow A$ and $g: A \rightarrow F(X)$ such that $fg = 1_A$ and $gf = 1_{F(X)}$.

Remarks.

- Clearly, if F is surjective, it is essentially surjective. But the reverse is not true.
- A functor is an <http://planetmath.org/EquivalenceOfCategories> equivalence iff it is <http://planetmath.org/FullFunctor> full, <http://planetmath.org/FaithfulFunctor> faithful and essentially surjective.
- **isomorphism-dense subcategory.** A full subcategory \mathcal{S} of a category \mathcal{C} is said to be *isomorphism-dense in \mathcal{C}* , if the inclusion functor $\mathcal{S} \hookrightarrow \mathcal{C}$ is essentially surjective. Since \mathcal{S} is full, the inclusion functor is full and faithful. As a result, \mathcal{S} is isomorphism-dense if the inclusion functor is an equivalence.