



Math for the people, by the people.

cochain complex

Canonical name	CochainComplex
Date of creation	2013-03-22 19:03:52
Last modified on	2013-03-22 19:03:52
Owner	rm50 (10146)
Last modified by	rm50 (10146)
Numerical id	5
Author	rm50 (10146)
Entry type	Definition
Classification	msc 18G35
Classification	msc 16E05
Defines	cocycle
Defines	coboundary

Let R be a ring. A sequence of <http://planetmath.org/ModuleR>-modules and homomorphisms

$$\dots \rightarrow A^{n-1} \xrightarrow{d_{n-1}} A^n \xrightarrow{d_n} A^{n+1} \xrightarrow{d_{n+1}} \dots$$

is said to be a *cochain complex* (or *R-complex*, or just *complex*) if each pair of adjacent homomorphisms (d_{n-1}, d_n) satisfies the relation $d_n \circ d_{n-1} = 0$. This is equivalent to saying that $\text{im } d_{n-1} \subseteq \ker d_n$. We often denote such a complex by (\mathcal{A}, d) , or simply \mathcal{A} .

Compare this to the notion of an exact sequence, which requires $\text{im } d_{n-1} = \ker d_n$. Compare also to the notion of a chain complex, in which the arrows go in the opposite direction.

The homomorphisms d_n in the chain complex are called *coboundary operators*, or *coboundary maps*. Elements of $\ker d_n$ are known as *cocycles*; elements of $\text{im } d_{n-1}$ as *coboundaries*.